

Bicyclopyrone
**Bicyclopyrone SL (A16003E) - *Salmonella Typhimurium* and
Escherichia Coli Reverse Mutation Assay**
Final Report

TEST GUIDELINE(S): OECD 471 (2020)

AUTHOR(S): Dr. Steffi Chang

COMPLETION DATE: 25 January 2021

PERFORMING LABORATORY: ICCR-Roßdorf GmbH
In den Leppsteinswiesen 19
64380 Rossdorf, Germany

LABORATORY PROJECT ID: Report Number: 2129100
Study Number: 2129100
Task Number: TK0539262

SPONSOR(S): Syngenta Ltd.
Jealott's Hill International Research Centre
Bracknell, Berkshire RG42 6EY, United Kingdom

STATEMENT OF DATA CONFIDENTIALITY CLAIMS

The Following Statement Applies To The United States of America:

STATEMENT OF NO DATA CONFIDENTIALITY CLAIMS UNDER SPECIFIED FIFRA PROVISIONS

No claim of confidentiality, on any basis whatsoever, is made for any information contained in this document. I acknowledge that information not designated as within the scope of FIFRA sec. 10(d)(1)(A), (B), or (C) and which pertains to a registered or previously registered pesticide is not entitled to confidential treatment and may be released to the public, subject to the provisions regarding disclosure to multinational entities under FIFRA 10(g).

Company: Syngenta Crop Protection, LLC
410 Swing Road
Post Office Box 18300
Greensboro, NC 27419-8300 USA

Submitter: _____ Date: _____

Syngenta is the owner of this information and data. Syngenta has submitted this material to the United States Environmental Protection Agency specifically under the provisions contained in FIFRA as amended and, hereby, consents to use and disclosure of this material by EPA according to FIFRA. In submitting this material to EPA according to method and format requirements contained in PR Notice 2011-3, we do not waive any protection or right involving this material that would have been claimed by the company if this material had not been submitted to the EPA, nor do we waive any protection or right provided under FIFRA Section 3 (concerning data exclusivity and data compensation) or FIFRA Section 10(g) (prohibiting disclosure to foreign and multinational pesticide companies or their agents).

GOOD LABORATORY PRACTICE COMPLIANCE STATEMENT

This study performed in the test facility of ICCR-Roßdorf GmbH, In den Leppsteinswiesen 19, 64380 Rossdorf, Germany was conducted in compliance with Good Laboratory Practice Regulations:

Chemikaliengesetz (Chemicals Act) of the Federal Republic of Germany, "Anhang 1" (Annex 1), in its currently valid version


OECD Principles of Good Laboratory Practice, (as revised in 1997), ENV/MC/CHEM(98)17

EC Commission Directive 2004/10/EC

These procedures are compatible with Good Laboratory Practice regulations specified by regulatory authorities throughout the European Community, the United States (EPA and FDA), and Japan (MHW, MAFF, and METI), and other countries that are signatories to the OECD Mutual Acceptance of Data Agreement.

There were no circumstances that may have affected the quality or integrity of the study.

Dr. Steffi Chang
Study Director Bacterial Systems


.....
Date: 25 January 2021

Performing Laboratory:
ICCR-Roßdorf GmbH
In den Leppsteinswiesen 19
64380 Rossdorf, Germany

To be completed for USA EPA submission only:

Representative of Submitter/Sponsor:

Date

Submitter/Sponsor: Syngenta Crop Protection, LLC
410 Swing Road
Post Office Box 18300
Greensboro, NC 27419-8300 USA

FLAGGING STATEMENT

This page is intentionally left blank. It will be replaced by an appropriate Flagging statement by the Sponsor.

QUALITY ASSURANCE STATEMENT

ICCR Study Number: 2129100
Test substance: Bicyclopyrone SL (A16003E)
Study director: Dr. Steffi Chang
Study Title: Bicyclopyrone SL (A16003E) -
Salmonella Typhimurium and
Escherichia Coli Reverse Mutation Assay

Study based activities at the Test Facility ICCR-Roßdorf GmbH were audited and inspected. The details of these audits and inspections are given below.

Type of Inspection	Date(s) of Inspection	Date Reporting to Study Director, Test Facility Management
Study Plan Verification	21 September 2020	21 September 2020
Process – based Test Item Preparation	21 & 24 September 2020	24 September 2020
Report Audit	09 November 2020	09 November 2020

General facilities and activities where this study was conducted were inspected on an annual basis and results are reported to the relevant responsible person and Management.

The statement is to confirm, that this report reflects the raw data.



H. Pilawa

Quality Assurance Auditor
ICCR-Roßdorf GmbH

25 January 2021

Date

PROJECT STAFF SIGNATURE

Study Director

Dr. Steffi Chang



.....
Date: 25 January 2021



GENERAL INFORMATION

Contributors

The following contributed to this report in the capacities indicated:

Name	Title
Dr. Steffi Chang	Study Director
Dr. Markus Schulz	Test Facility Management
Frauke Hermann	Head of Quality Assurance Unit
Carolina Vaccari	Syngenta Study Manager

Study Dates

Study initiation date:	22 September 2020
Experimental start date:	01 October 2020
Experimental completion date:	21 October 2020

Deviations from the Guidelines

None

Retention of Samples

None

Performing Laboratory Test Substance Reference Number

S 2111411

Other

ICCR-Roßdorf GmbH will archive:

Records and documentation relating to this study will be maintained in the archives of ICCR-Roßdorf GmbH for a period of 4 years from the date on which the Study Director signs the final report. This will include electronic and paper raw data, and report that support the reconstruction of the study.

At termination of the aforementioned period, the records and documentation will be transferred to the GLP compliant Archive of Rhenus Archiv Services GmbH, Frankfurt am Main for further archiving up to a total archiving period of 15 years.

A sample of the test item will not be archived.

ICCR Roßdorf GmbH will retain in its archive a copy of the study plan and final report, and any amendments indefinitely.

Deviations from the study plan

There were no deviations (unplanned changes) from the study plan.

Distribution of the report

Sponsor	2 × electronic copy (1 × pdf-file, 1 × Word-file)
Study Director	1 × (original)

TABLE OF CONTENTS

STATEMENT OF DATA CONFIDENTIALITY CLAIMS	2
GOOD LABORATORY PRACTICE COMPLIANCE STATEMENT	3
FLAGGING STATEMENT	4
QUALITY ASSURANCE STATEMENT	5
PROJECT STAFF SIGNATURE	6
GENERAL INFORMATION	7
TABLE OF CONTENTS	9
1.0 EXECUTIVE SUMMARY	11
1.1 Study Design	11
1.2 Results	11
1.3 Conclusion.....	11
2.0 INTRODUCTION	12
2.1 Purpose.....	12
2.2 Test Guideline(s).....	12
3.0 MATERIALS AND METHODS	13
3.1 Test Substance.....	13
3.2 Controls	14
3.2.1 Negative controls	14
3.2.2 Positive control substances	14
3.3 Experimental Design.....	15
3.3.1 Characterisation of the <i>Salmonella typhimurium</i> and <i>E. coli</i> strains	15
3.3.2 Storage.....	16
3.3.3 Precultures.....	16
3.3.4 Selective agar	16
3.3.5 Overlay agar	16
3.4 Mammalian Microsomal Fraction S9 Mix.....	16
3.4.1 S9 (Preparation by ICCR-Roßdorf GmbH).....	16
3.4.2 S9 mix	17
3.5 Pre-Experiment for Cytotoxicity.....	17
3.6 Concentration Selection	18
3.7 Experimental Performance.....	18
3.8 Data Evaluation	18

3.8.1	Data recording	18
3.8.2	Acceptability of the assay	19
3.8.3	Evaluation of results.....	19
3.8.4	Biometry.....	19
3.9	Major Computerized System.....	19
4.0	RESULTS AND DISCUSSION	20
5.0	CONCLUSIONS	20
6.0	REFERENCES	21
TABLES SECTION		22
TABLE 1	Summary of Results Pre-Experiment/Experiment I.....	23
TABLE 2	Summary of Results Experiment II.....	24
TABLE 3	Pre-Experiment and Experiment I: 2129100 VV Plate Incorporation Without Metabolic Activation.....	25
TABLE 4	Pre-Experiment and Experiment I: 2129100 VV Plate Incorporation With Metabolic Activation.....	27
TABLE 5	Experiment II: 2129100 HV2 Pre Incubation Without Metabolic Activation.....	29
TABLE 6	Experiment II: 2129100 HV2 Pre Incubation With Metabolic Activation.....	31
APPENDICES SECTION		33
APPENDIX 1	Historical Control Data	34
APPENDIX 2	Copy of GLP Certificate	35
APPENDIX 3	Certificate of S9	36
APPENDIX 4	Certificate of Analysis.....	37

1.0 EXECUTIVE SUMMARY

1.1 Study Design

This study was performed to investigate the potential of Bicyclopyrone SL (A16003E) to induce gene mutations in the plate incorporation test (Experiment I) and the pre-incubation test (Experiment II) using the *Salmonella typhimurium* (*S. typhimurium*) strains TA1535, TA1537, TA98, and TA100, and the *Escherichia coli* (*E. coli*) strains WP2 *uvrA* (pKM101) and WP2 (pKM101).

1.2 Results

The plates incubated with the test item showed normal background growth up to the maximal concentration of 5000 µg/plate with and without S9 mix in all strains used.

No cytotoxic effects, evident as a reduction in the number of revertants (below the indication factor of 0.5), occurred in all strains with and without metabolic activation.

No relevant increase in revertant colony numbers of any of the six tester strains was observed following treatment with Bicyclopyrone SL (A16003E) at any concentration level, neither in the presence nor absence of metabolic activation (S9 mix). There was also no observed tendency of higher mutation rates with increasing concentrations in the range below the generally acknowledged border of biological relevance.

Appropriate reference mutagens were used as positive controls, which showed a distinct increase of induced revertant colonies consistent with the laboratory's historical control data demonstrated the sensitivity of the test system and the efficacy of the S9 mix. Each batch of S9 was also tested with 2 pro-mutagens, benzo(a)pyrene and 2-aminoanthracene.

1.3 Conclusion

In conclusion, it can be stated that during the described mutagenicity tests and under the experimental conditions reported, Bicyclopyrone SL (A16003E) did not induce gene mutations by base pair changes or frameshifts in the genome of the strains used. Therefore, Bicyclopyrone SL (A16003E) is considered to be non-mutagenic in the *Salmonella typhimurium* and *Escherichia coli* reverse mutation assay.

2.0 INTRODUCTION

2.1 Purpose

These experiments were performed to assess the potential of the test substance to induce gene mutations by means of the *S. typhimurium* and *E. coli* reverse mutation assay. Experiment I was performed as a plate incorporation assay. Since a negative result was obtained in this experiment, Experiment II was performed as a pre-incubation assay.

The most widely used assays for detecting gene mutations are those using bacteria (1). They are relatively simple and rapid to perform, and give reliable data on the ability of an agent to interact with DNA and produce mutations.

Reverse mutation assays determine the frequency with which an agent reverses or suppresses the effect of the forward mutation. The genetic target presented to an agent is therefore small, specific and selective. Several bacterial strains, or a single strain with multiple markers are necessary to assure reliable detection of mutagens that may be specific to one tester strain or locus. The reversion of bacteria from growth-dependence on a particular amino acid to growth in the absence of that amino acid (reversion from auxotrophy to prototrophy) is the most widely used marker.

The *S. typhimurium* histidine (his) and the *E. coli* tryptophan (trp) reversion system measures his⁻ → his⁺ and trp⁻ → trp⁺ reversions, respectively. The *S. typhimurium* and *E. coli* strains are constructed to differentiate between base pair (TA1535, TA100, WP2 *uvrA* (pKM101), and WP2 (pKM101)) and frameshift (TA1537, TA98) mutations.

According to the direct plate incorporation and pre-incubation method the bacteria are exposed to the test substance with and without metabolic activation and plated on selective medium. After a suitable period of incubation, revertant colonies are counted.

To establish a concentration response effect at least six concentrations with adequately spaced intervals were tested. The maximum concentration was 5000 µg/plate.

To validate the test, reference mutagens were tested in parallel to the test substance.

2.2 Test Guideline(s)

This study followed the procedures indicated by the following internationally accepted guideline and recommendations:

“Ninth Addendum to OECD Guidelines for Testing of Chemicals”, Section 4, No. 471: “Bacterial Reverse Mutation Test”, corrected June 26, 2020

3.0 MATERIALS AND METHODS

3.1 Test Substance

Information as provided by the Sponsor.

Identification:	Bicyclopyrone SL (A16003E)
Batch:	1099852
Concentration of Bicyclopyrone SL (A16003E):	18.5%
Appearance:	Brown transparent liquid
Recertification Date:	29 August 2022
Storage Conditions:	At room temperature
Stability in Solvent:	Not indicated by the Sponsor

The test substance concentrations were not adjusted for the content of Bicyclopyrone SL (A16003E).

On the day of the experiment (immediately before use), the test substance was dissolved in dimethylsulfoxide (DMSO, purity > 99%). The solvent was chosen as the more suitable solvent compared to water, according to its solubilisation properties and its relative non-toxicity to the bacteria (2).

All formulations were prepared freshly before treatment and used within two hours of preparation. The formulation was assumed to be stable for this period unless specified otherwise by the Sponsor.

3.2 Controls

3.2.1 Negative controls

Concurrent untreated and solvent controls were performed.

3.2.2 Positive control substances

Without metabolic activation

Strains: TA1535, TA100
Name: Sodium azide, (NaN₃)
Supplier: SERVA, 69042 Heidelberg, Germany
Batch No.: 150564
Purity: ≥ 99%
Dissolved in: Deionised water
Concentration: 10 µg/plate

Strains: TA1537, TA98
Name: 4-nitro-*o*-phenylene-diamine, (4-NOPD)
Supplier: Sigma-Aldrich, 82024 Taufkirchen, Germany
Batch No.: MKBM 5257V
Purity: ≥ 98%
Dissolved in: DMSO (purity >99 %, Fisher Leics LE11 5RG, United Kingdom)
Concentration: 10 µg/plate in strain TA 98, 50 µg/plate in strain TA 1537

Strains: WP2 *uvrA* (pKM101), WP2 (pKM101)
Name: Methyl methane sulfonate, (MMS)
Supplier: Sigma-Aldrich, 82024 Taufkirchen, Germany
Batch No.: MKCG 1346
Purity: ≥ 99%
Dissolved in: Deionised water
Concentration: 2.0 µL/plate

With metabolic activation

Strains: TA1535, TA1537, TA98, TA100, WP2 *uvrA* (pKM101), WP2 (pKM 101)
Name: 2-aminoanthracene, (2-AA)
Supplier: Sigma-Aldrich, 82024 Taufkirchen, Germany
Batch No.: STBG 0630V
Purity: ≥ 96%
Dissolved in: DMSO (purity > 99 %, Fisher Leics LE11 5RG, United Kingdom)
Concentration: 2.5 µg/plate (TA1535, TA1537, TA98, TA100),
10 µg/plate (WP2 *uvrA* (pKM101), WP2 (pKM101))

The stability of the positive control substances in solution is unknown but a mutagenic response in the expected range is sufficient evidence of biological activity.

3.3 Experimental Design

3.3.1 Characterisation of the *Salmonella typhimurium* and *E. coli* strains

The histidine dependent strains are derived from *S. typhimurium* strain LT2 through mutations in the histidine locus. Additionally, due to the "deep rough" (*rfa*⁻) mutation they possess a faulty lipopolysaccharide envelope which enables substances to penetrate the cell wall more easily. A further mutation causes a reduction in the activity of an excision repair system. The last alteration includes mutational processes in the nitrate reductase and biotin genes produced in a UV-sensitive area of the gene named *uvrB*⁻. In the strains TA98 and TA100 the R-factor plasmid pKM101 carries the ampicillin resistance marker (3).

Strain WP2 (4) and its derivatives all carry the same defect in one of the genes for tryptophan biosynthesis. Tryptophan-independent (*Trp*⁺) mutants (revertants) can arise either by a base change at the site of the original alteration or by a base change elsewhere in the chromosome so that the original defect is suppressed. This second possibility can occur in several different ways so that the system seems capable of detecting all types of mutagen which substitute one base for another. Additionally, the *uvrA* derivative is deficient in the DNA repair process (excisable repair damage). Such a repair-deficient strain may be more readily mutated by agents. The *E. coli* strains WP2 *uvrA* (pKM101) and WP2 (pKM101) are constructed by introduction of the R-factor plasmid pKM101.

When summarized, the mutations of the *S. typhimurium* and *E. coli* strains used in this study can be described as follows:

Strains	Genotype	Type of mutations indicated
<i>Salmonella typhimurium</i>		
TA1537	<i>his C 3076; rfa</i> ⁻ ; <i>uvrB</i> ⁻	frame shift mutations
TA98	<i>his D 3052; rfa</i> ⁻ ; <i>uvrB</i> ⁻ ; R-factor	" "
TA1535	<i>his G 46; rfa</i> ⁻ ; <i>uvrB</i> ⁻	base-pair substitutions
TA100	<i>his G 46; rfa</i> ⁻ ; <i>uvrB</i> ⁻ ; R-factor	" "
<i>Escherichia coli</i>		
WP2 <i>uvrA</i> (pKM101)	<i>trp E 56 uvrA</i> ⁻ ; R-factor	base-pair substitutions and others
WP2 (pKM101)	<i>trp E 56</i> ; R-factor	" "

Regular checking of the properties of the *S. typhimurium* and *E. coli* strains regarding the membrane permeability and ampicillin resistance; UV sensitivity, and amino acid requirement as well as normal spontaneous mutation rates is performed by ICCR-Roßdorf GmbH according to Ames *et al.* (5), Maron and Ames (3), and Mortelmans and Riccio (7). In this way it is ensured that the experimental conditions set down by Ames are fulfilled.

The bacterial strains TA1535, TA1537, TA98, TA100, WP2 *uvrA* (pKM101), and WP2 (pKM101) were obtained from Trinova Biochem GmbH (35394 Gießen, Germany).

3.3.2 Storage

The strain cultures were stored as stock cultures in ampoules with nutrient broth + 5 % DMSO (Fisher Leics, LE11 5RG, United Kingdom) in liquid nitrogen.

3.3.3 Precultures

The thawed bacterial suspension was transferred into 250 mL Erlenmeyer flasks containing nutrient medium (50 mL). A solution of ampicillin (50 µL, 25 µg/mL) was added to the strains TA98, TA100, WP2 *uvrA* (pKM101), and WP2 (pKM101). This nutrient medium contains per liter:

8 g Nutrient Broth (MERCK, 64293 Darmstadt, Germany)

5 g NaCl (MERCK, 64293 Darmstadt, Germany)

The bacterial cultures were incubated in a shaking water bath for 4 hours at 37 °C. The optical density of the bacteria was determined by absorption measurement and the obtained values indicated that the bacteria were harvested at the late exponential or early stationary phase (10^8 - 10^9 cells/mL).

3.3.4 Selective agar

Plates with selective agar (without Histidine/Tryptophan) were used.

3.3.5 Overlay agar

The overlay agar contained per litre:

for *Salmonella* strains:

7.0 g Agar Agar*

6.0 g NaCl*

10.5 mg L-Histidine×HCl×H₂O*

12.2 mg Biotin*

for *Escherichia coli* strains:

7.0 g Agar Agar*

6.0 g NaCl*

10.2 mg Tryptophan*

* (MERCK, 64293 Darmstadt, Germany)

Sterilisations were performed at 121 °C in an autoclave.

3.4 Mammalian Microsomal Fraction S9 Mix

The bacteria used in this assay do not possess the enzyme systems which, in mammals, are known to convert promutagens into active DNA damaging metabolites. In order to overcome this major drawback an exogenous metabolic system is added in the form of mammalian microsome enzyme activation mixture.

3.4.1 S9 (Preparation by ICCR-Roßdorf GmbH)

Phenobarbital/β-naphthoflavone induced rat liver S9 was used as the metabolic activation system. The S9 was prepared from male Wistar rats (RjHan:WI; weight approx. 220 – 320 g,

Janvier Labs, 53941 Saint-Berthevin Cedex, France) induced by peroral administration of 80 mg/kg b.w. phenobarbital (Sigma-Aldrich Chemie GmbH, 82024 Taufkirchen, Germany) and by peroral administrations of β -naphthoflavone (Acros Organics, 2440 Geel, Belgium) each, on three consecutive days. The livers were prepared 24 hours after the last treatment. The S9 fractions were produced by dilution of the liver homogenate with a KCl solution (1+3 parts) followed by centrifugation at 9000 g. Aliquots of the supernatant were frozen and stored in ampoules at $-80\text{ }^{\circ}\text{C}$. Small numbers of the ampoules can be kept at $-20\text{ }^{\circ}\text{C}$ for up to one week. Each batch of S9 mix is routinely tested with 2-aminoanthracene as well as benzo[a]pyrene (Appendix 3).

The protein concentration in the S9 preparation was 33.0 mg/mL (lot no. 030920K) in both experiments.

3.4.2 S9 mix

Before the experiment an appropriate quantity of S9 supernatant was thawed and mixed with S9 cofactor solution. The amount of S9 supernatant was 10% v/v in the S9 mix. Cofactors were added to the S9 mix to reach the following concentrations in the S9 mix:

8 mM MgCl_2
33 mM KCl
5 mM Glucose-6-phosphate
4 mM NADP

in 100 mM sodium-ortho-phosphate-buffer, pH 7.4.

During the experiment the S9 mix was stored in an ice bath. The S9 mix preparation was performed according to Ames *et al.*(5).

3.5 Pre-Experiment for Cytotoxicity

To evaluate the cytotoxicity of the test substance a pre-experiment was performed with all strains. Eight concentrations were tested for cytotoxicity and mutation induction each with three replicate plates. The experimental conditions in this pre-experiment are described in section 3.7 (plate incorporation test).

Cytotoxicity of the test substance results in a reduction in the number of spontaneous revertants (below a factor of 0.5) or a clearing of the bacterial background lawn.

The pre-experiment is reported as the Main Experiment I since the criteria mentioned in Section 3.8.2 Acceptability of the Assay were met.

3.6 Concentration Selection

In the pre-experiment the concentration range of the test substance was 3 - 5000 µg/plate. The pre-experiment is reported as Experiment I. Since no cytotoxic effects were observed 5000 µg/plate was chosen as the maximal concentration in Experiment II.

The concentration range included two logarithmic decades. The following concentrations were tested in experiment II:

33; 100; 333; 1000; 2500; and 5000 µg/plate

3.7 Experimental Performance

For each strain and concentration including the controls, three plates were used.

The following materials were mixed in a test tube and poured onto the selective agar plates:

- 100 µL Test solution at each concentration, solvent (negative control) or reference mutagen solution (positive control),
- 500 µL S9 mix (for test with metabolic activation) or S9 mix substitution buffer* (for test without metabolic activation),
- 100 µL Bacteria suspension (cf. test system, pre-culture of the strains; OD = 1.0 - 1.2; wavelength = 500 nm; approx. 8×10^8 cells/mL),
- 2000 µL Overlay agar

For the pre-incubation method test solution (100 µL) (solvent or reference mutagen solution (positive control)), S9 mix / S9 mix substitution buffer* (500 µL) and bacteria suspension (100 µL) were mixed in a test tube and incubated at $37 \text{ C} \pm 1.5^\circ \text{ C}$ for 60 minutes. After pre-incubation overlay agar (2.0 mL, 45° C) was added to each tube. The mixture was poured on selective agar plates.

After solidification the plates were incubated upside down for 72 hours at $37 \text{ C} \pm 1.5^\circ \text{ C}$ in the dark, plates were then stored at 4° C until counted (6).

In parallel to each test a sterile control of the test substance was performed and documented in the raw data. Therefore, stock solution (100 µL) and S9 mix / S9 mix substitution buffer* (500 µL) were mixed with overlay agar (2.0 mL) and poured on minimal agar plates.

* Substitution buffer: 7 parts of the 100 mM sodium-ortho-phosphate-buffer pH 7.4 with 3 parts of KCl solution 0.15 M

3.8 Data Evaluation

3.8.1 Data recording

The colonies were counted using a Petri Viewer with the software program Ames Study Manager (see section 3.9, Major computerized systems). The evaluation unit was connected

to a PC with printer to print out the individual values, the means from the plates for each concentration together with standard deviations and enhancement factors as compared to the spontaneous reversion rates (see tables of results). The print outs are kept with the raw data.

3.8.2 Acceptability of the assay

The *Salmonella typhimurium* and *Escherichia coli* reverse mutation assay is considered acceptable if it meets the following criteria:

- regular background growth in the negative and solvent control
- the spontaneous reversion rates in the negative and solvent control are in the range of the historical data
- the positive control substances should produce an increase in mutant colony frequencies of at least 2-fold concurrent control
- a minimum of five analysable concentrations should be present with at least four showing no signs of toxic effects, evident as a reduction in the number of revertants below the indication factor of 0.5.

3.8.3 Evaluation of results

A test substance is considered as a mutagen if a biologically relevant increase in the number of revertants of twice or above the spontaneous mutation rate of the corresponding solvent control is observed (1).

A concentration dependent increase is considered biologically relevant if the threshold is reached or exceeded at more than one concentration (6).

An increase of revertant colonies equal or above the threshold at only one concentration is judged as biologically relevant if reproduced in an independent second experiment.

A concentration dependent increase in the number of revertant colonies below the threshold is regarded as an indication of a mutagenic potential if reproduced in an independent second experiment. However, whenever the colony counts remain within the historical range of negative and solvent controls, such an increase is not considered biologically relevant.

3.8.4 Biometry

According to the OECD guideline 471, a statistical analysis of the data is not mandatory.

3.9 Major Computerized System

Petri Viewer Sorcerer Colony Counter 3.0 (Instem, Suffolk IP33 3TA, UK) with the software program Ames Study Manager (v1.24) and Ames Archive Manager (v1.01).

4.0 RESULTS AND DISCUSSION

The test substance, Bicyclopyrone SL (A16003E), was assessed for its potential to induce gene mutations in the plate incorporation test (Experiment I) and the pre-incubation test (Experiment II) using *S. typhimurium* strains TA1535, TA1537, TA98, and TA100, and the *E. coli* strains WP2 (pKM101) and WP2 *uvrA* (pKM101).

In the pre-experiment the concentration range of the test substance was 3 - 5000 µg/plate. The pre-experiment is reported as Experiment I. Since no cytotoxic effects were observed 5000 µg/plate was chosen as the maximal concentration in Experiment II.

The assay was performed with and without liver microsomal activation. Each concentration, including the controls, was tested in triplicate. The concentration range included two logarithmic decades. The test substance was tested at the following concentrations:
Pre-Experiment/Experiment I: 3; 10; 33; 100; 333; 1000; 2500; and 5000 µg/plate
Experiment II: 33; 100; 333; 1000; 2500; and 5000 µg/plate

No precipitation of the test item occurred up to the highest investigated dose.

The plates incubated with the test item showed normal background growth up to the maximal dose of 5000 µg/plate with and without S9 mix in all strains used.

No cytotoxic effects, evident as a reduction in the number of revertants (below the indication factor of 0.5), occurred in the test groups with and without metabolic activation.

No substantial increase in revertant colony numbers in any of the six tester strains was observed following treatment with Bicyclopyrone SL (A16003E) at any concentration level, neither in the presence nor absence of metabolic activation (S9 mix). There was also no tendency of higher mutation rates with increasing concentrations in the range below the generally acknowledged border of biological relevance.

Appropriate reference mutagens were used as positive controls. They showed a distinct increase in induced revertant colonies.

5.0 CONCLUSIONS

In conclusion, it can be stated that during the described mutagenicity tests and under the experimental conditions reported, Bicyclopyrone SL (A16003E) did not induce gene mutations by base pair changes or frameshifts in the genome of the strains used.

Therefore, Bicyclopyrone SL (A16003E) is considered to be non-mutagenic in the *Salmonella typhimurium* and *Escherichia coli* reverse mutation assay.

6.0 REFERENCES

1. Hollstein, M., J. McCann, F.A. Angelosanto, and W.W. Nichols (1979)
Short-term tests for carcinogens and mutagens
Mutation Res. 65, 133-226
2. Maron, D.M., J. Katzenellenbogen, and B.N. Ames (1981)
Compatibility of organic solvents with the Salmonella/Microsome Test
Mutation Res. 88, 343-350
3. Maron, D.M. and B.N. Ames (1983)
Revised methods for the Salmonella mutagenicity test
Mutation Res. 113, 173-215
4. Green, M.H.L. and W.J. Muriel (1976)
Mutagen Testing Using TRP⁺ Reversion in Escherichia Coli
Mutation Res. 38, 3-32
5. Ames, B.N., J. McCann, and E. Yamasaki (1977)
Methods for detecting carcinogens and mutagens with the Salmonella/mammalian
microsome mutagenicity test
In: B.J. Kilbey et al. (Eds.) "Handbook of Mutagenicity Test Procedures" Elsevier,
Amsterdam, 1-17
6. de Serres, F.J. and M.D. Shelby (1979)
Recommendations on data production and analysis using the Salmonella/microsome
mutagenicity assay
Mutation Res. 64, 159-165
7. Mortelmans, K. and E.S. Riccio (2000)
The bacterial tryptophan reverse mutation assay with Escherichia coli WP2
Mutation Res. 455, 61-69

TABLES SECTION

TABLE 1 Summary of Results Pre-Experiment/Experiment I

Study Name: 2129100
 Experiment: 2129100 VV Plate
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 01.10.2020
 Date Counted: 08.10.2020

Metabolic Activation	Test Group	Concentration (per plate)	Revertant Colony Counts (Mean ±SD)					
			TA 1535	TA 1537	TA 98	TA 100	WP2 pKM101	WP2 uvrA pKM101
Without Activation	DMSO Untreated		9 ± 4	9 ± 3	35 ± 8	110 ± 1	213 ± 16	284 ± 6
	Bicyclopyrone SL (A16003E)	3 µg	9 ± 2	10 ± 4	44 ± 6	94 ± 9	217 ± 12	275 ± 22
		10 µg	8 ± 3	11 ± 1	41 ± 4	106 ± 1	195 ± 19	270 ± 18
		33 µg	10 ± 0	10 ± 1	36 ± 6	113 ± 26	221 ± 35	282 ± 11
		100 µg	9 ± 2	12 ± 3	40 ± 10	96 ± 12	215 ± 9	304 ± 16
		333 µg	12 ± 2	9 ± 2	41 ± 6	98 ± 8	197 ± 7	310 ± 13
		1000 µg	10 ± 1	10 ± 1	44 ± 11	94 ± 6	199 ± 19	258 ± 20
		2500 µg	8 ± 3	8 ± 1	35 ± 6	103 ± 10	205 ± 22	265 ± 3
		5000 µg	9 ± 3	8 ± 1	35 ± 3	97 ± 9	199 ± 13	285 ± 25
	NaN3	10 µg	999 ± 20			1396 ± 53		
	4-NOPD	10 µg			583 ± 20			
	4-NOPD	50 µg		91 ± 10				
	MMS	2.0 µL					2712 ± 183	2497 ± 219
	With Activation	DMSO Untreated		11 ± 0	13 ± 4	50 ± 2	76 ± 9	207 ± 24
Bicyclopyrone SL (A16003E)		3 µg	10 ± 1	12 ± 2	50 ± 2	82 ± 12	234 ± 6	330 ± 12
		10 µg	11 ± 4	14 ± 2	53 ± 7	80 ± 4	232 ± 12	320 ± 10
		33 µg	9 ± 2	11 ± 1	47 ± 5	85 ± 7	213 ± 17	299 ± 18
		100 µg	11 ± 1	12 ± 3	50 ± 3	89 ± 6	183 ± 8	288 ± 12
		333 µg	11 ± 1	13 ± 1	47 ± 6	91 ± 9	216 ± 6	299 ± 23
		1000 µg	13 ± 2	13 ± 4	42 ± 6	77 ± 17	189 ± 15	311 ± 29
		2500 µg	9 ± 3	12 ± 3	44 ± 3	85 ± 9	211 ± 12	284 ± 23
5000 µg		9 ± 3	8 ± 2	35 ± 9	86 ± 8	206 ± 21	303 ± 25	
2-AA		2.5 µg	12 ± 2	10 ± 1	47 ± 10	93 ± 16	207 ± 39	301 ± 10
2-AA	10.0 µg	200 ± 24	371 ± 37	2141 ± 424	2438 ± 243	976 ± 31	1567 ± 118	

Key to Positive Controls

NaN3	sodium azide
2-AA	2-aminoanthracene
4-NOPD	4-nitro-o-phenylene-diamine
MMS	methyl methane sulfonate

TABLE 2 Summary of Results Experiment II

Study Name: 2129100
 Experiment: 2129100 HV2 Pre
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 15.10.2020
 Date Counted: 21.10.2020

Metabolic Activation	Test Group	Concentration (per plate)	Revertant Colony Counts (Mean ±SD)						
			TA 1535	TA 1537	TA 98	TA 100	WP2 pKM101	WP2 uvrA pKM101	
Without Activation	DMSO Untreated		7 ± 1	14 ± 3	33 ± 3	93 ± 4	246 ± 25	326 ± 19	
	Bicyclopyrone SL (A16003E)	33 µg	10 ± 2	10 ± 1	22 ± 2	101 ± 11	268 ± 13	324 ± 8	
		100 µg	7 ± 1	13 ± 4	30 ± 7	102 ± 1	253 ± 9	289 ± 25	
		333 µg	10 ± 2	13 ± 2	29 ± 8	82 ± 14	232 ± 12	311 ± 10	
		1000 µg	9 ± 3	13 ± 2	24 ± 7	86 ± 8	241 ± 27	316 ± 8	
		2500 µg	7 ± 2	11 ± 1	22 ± 5	82 ± 12	236 ± 31	299 ± 19	
		5000 µg	6 ± 2	14 ± 2	26 ± 8	76 ± 7	218 ± 22	266 ± 14	
	NaN3	10 µg	1047 ± 29			1578 ± 137			
	4-NOPD	10 µg			702 ± 56				
	4-NOPD MMS	50 µg 2.0 µL		105 ± 12			2830 ± 24	2797 ± 81	
	With Activation	DMSO Untreated		10 ± 3	13 ± 3	41 ± 4	87 ± 4	271 ± 30	323 ± 9
		Bicyclopyrone SL (A16003E)	33 µg	11 ± 2	17 ± 2	43 ± 10	112 ± 7	290 ± 3	372 ± 15
100 µg			12 ± 3	13 ± 2	40 ± 2	95 ± 10	267 ± 14	348 ± 27	
333 µg			10 ± 2	12 ± 2	38 ± 6	92 ± 8	241 ± 24	332 ± 32	
1000 µg			9 ± 1	15 ± 0	35 ± 12	100 ± 7	242 ± 11	325 ± 17	
2500 µg			9 ± 3	13 ± 3	39 ± 7	97 ± 10	260 ± 16	332 ± 22	
5000 µg			11 ± 3	14 ± 5	37 ± 9	91 ± 10	250 ± 34	330 ± 18	
2-AA		2.5 µg	12 ± 3	12 ± 1	36 ± 12	107 ± 6	266 ± 10	326 ± 16	
2-AA		10.0 µg	259 ± 18	402 ± 44	3134 ± 159	3501 ± 248	941 ± 12	1860 ± 42	

Key to Positive Controls

NaN3	sodium azide
2-AA	2-aminoanthracene
4-NOPD	4-nitro-o-phenylene-diamine
MMS	methyl methane sulfonate

TABLE 3 Pre-Experiment and Experiment I: 2129100 VV Plate Incorporation Without Metabolic Activation

Study Name: 2129100
 Experiment: 2129100 VV Plate
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 01.10.2020
 Date Counted: 08.10.2020

Without metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
TA 1535	Bicyclopyrone SL (A16003E)	3 µg	9.0	1.7	1.0	7, 10, 10
		10 µg	8.3	3.2	0.9	12, 7, 6
		33 µg	10.0	0.0	1.1	10, 10, 10
		100 µg	9.0	1.7	1.0	10, 7, 10
		333 µg	12.0	1.7	1.3	11, 11, 14
		1000 µg	10.0	1.0	1.1	9, 11, 10
		2500 µg	8.0	2.6	0.9	6, 11, 7
		5000 µg	9.3	2.9	1.0	11, 11, 6
	DMSO Untreated		9.3	4.0		7, 7, 14
		12.7	3.8		11, 10, 17	
TA 1537	Bicyclopyrone SL (A16003E)	3 µg	10.3	3.5	1.1	10, 14, 7
		10 µg	10.7	0.6	1.1	11, 10, 11
		33 µg	10.3	0.6	1.1	10, 11, 10
		100 µg	12.0	3.0	1.3	12, 15, 9
		333 µg	8.7	1.5	0.9	7, 10, 9
		1000 µg	10.3	0.6	1.1	10, 11, 10
		2500 µg	7.7	1.2	0.8	7, 7, 9
		5000 µg	8.3	1.2	0.9	9, 7, 9
	DMSO Untreated		9.3	2.5		12, 9, 7
		8.3	3.2		7, 6, 12	
TA 98	Bicyclopyrone SL (A16003E)	3 µg	44.0	6.1	1.3	37, 47, 48
		10 µg	41.0	4.4	1.2	36, 44, 43
		33 µg	36.3	5.5	1.0	42, 31, 36
		100 µg	40.3	9.7	1.2	51, 32, 38
		333 µg	41.3	5.5	1.2	47, 41, 36
		1000 µg	43.7	11.0	1.2	40, 56, 35
		2500 µg	34.7	6.4	1.0	42, 31, 31
		5000 µg	35.0	3.0	1.0	32, 38, 35
	DMSO Untreated		35.0	7.5		36, 27, 42
		42.0	4.6		38, 41, 47	
TA 100	Bicyclopyrone SL (A16003E)	3 µg	94.3	9.1	0.9	86, 104, 93
		10 µg	105.7	0.6	1.0	105, 106, 106
		33 µg	113.0	25.7	1.0	104, 93, 142
		100 µg	96.0	12.1	0.9	85, 94, 109
		333 µg	98.3	8.1	0.9	89, 104, 102
		1000 µg	94.0	6.0	0.9	94, 100, 88
		2500 µg	103.3	10.1	0.9	94, 102, 114
		5000 µg	96.7	8.6	0.9	89, 106, 95
	DMSO Untreated		110.0	1.0		109, 110, 111
		88.3	7.4		94, 80, 91	

Study Name: 2129100
 Experiment: 2129100 VV Plate
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 01.10.2020
 Date Counted: 08.10.2020

Without metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
WP2 pKM101	Bicyclopyrone SL (A16003E)	3 µg	216.7	11.5	1.0	228, 205, 217
		10 µg	195.0	19.1	0.9	173, 205, 207
		33 µg	221.0	34.8	1.0	216, 189, 258
		100 µg	215.3	8.5	1.0	209, 212, 225
		333 µg	197.3	6.7	0.9	199, 203, 190
		1000 µg	199.0	19.0	0.9	204, 215, 178
		2500 µg	204.7	22.1	1.0	230, 195, 189
		5000 µg	199.0	13.1	0.9	185, 201, 211
	DMSO		213.3	15.9		224, 195, 221
	Untreated		235.3	7.5		228, 235, 243
WP2 uvrA pKM101	Bicyclopyrone SL (A16003E)	3 µg	275.3	22.3	1.0	264, 261, 301
		10 µg	270.3	18.3	1.0	264, 256, 291
		33 µg	282.0	11.1	1.0	272, 294, 280
		100 µg	304.3	16.1	1.1	286, 316, 311
		333 µg	310.3	12.9	1.1	325, 301, 305
		1000 µg	258.0	20.1	0.9	272, 235, 267
		2500 µg	264.7	3.1	0.9	262, 268, 264
		5000 µg	285.3	25.1	1.0	294, 257, 305
	DMSO		284.3	5.5		290, 279, 284
	Untreated		313.7	16.8		303, 333, 305
TA 1535	NaN3	10 µg	998.7	19.5	107.0	990, 985, 1021
TA 1537	4-NOPD	50 µg	90.7	10.3	9.7	82, 102, 88
TA 98	4-NOPD	10 µg	583.3	19.9	16.7	569, 575, 606
TA 100	NaN3	10 µg	1395.7	53.5	12.7	1416, 1436, 1335
WP2 pKM101	MMS	2.0 µL	2712.0	183.0	12.7	2576, 2640, 2920
WP2 uvrA pKM101	MMS	2.0 µL	2496.7	219.5	8.8	2309, 2443, 2738

Key to Positive Controls

NaN3	sodium azide
4-NOPD	4-nitro-o-phenylene-diamine
MMS	methyl methane sulfonate

TABLE 4 Pre-Experiment and Experiment I: 2129100 VV Plate Incorporation With Metabolic Activation

Study Name: 2129100
 Experiment: 2129100 VV Plate
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 01.10.2020
 Date Counted: 08.10.2020

With metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
TA 1535	Bicyclopyrone SL (A16003E)	3 µg	10.7	3.5	1.0	11, 7, 14
		10 µg	8.7	2.3	0.8	10, 10, 6
		33 µg	11.3	1.2	1.0	10, 12, 12
		100 µg	11.0	1.0	1.0	11, 12, 10
		333 µg	12.7	2.1	1.2	11, 12, 15
		1000 µg	9.3	3.1	0.8	6, 10, 12
		2500 µg	9.3	2.5	0.8	7, 9, 12
		5000 µg	12.3	2.3	1.1	15, 11, 11
	DMSO		11.0	0.0		11, 11, 11
	Untreated		10.0	1.0		9, 11, 10
TA 1537	Bicyclopyrone SL (A16003E)	3 µg	14.0	2.0	1.1	12, 16, 14
		10 µg	11.0	1.0	0.8	11, 10, 12
		33 µg	12.3	2.5	0.9	12, 10, 15
		100 µg	13.3	1.2	1.0	14, 14, 12
		333 µg	13.0	3.6	1.0	9, 16, 14
		1000 µg	12.0	3.5	0.9	10, 10, 16
		2500 µg	8.3	2.1	0.6	9, 6, 10
		5000 µg	10.3	1.2	0.8	11, 11, 9
	DMSO		13.0	3.6		16, 14, 9
	Untreated		12.3	1.5		12, 14, 11
TA 98	Bicyclopyrone SL (A16003E)	3 µg	52.7	6.5	1.0	53, 46, 59
		10 µg	47.3	4.5	0.9	52, 47, 43
		33 µg	50.0	2.6	1.0	47, 51, 52
		100 µg	46.7	6.1	0.9	52, 48, 40
		333 µg	42.3	5.9	0.8	49, 38, 40
		1000 µg	44.3	2.9	0.9	41, 46, 46
		2500 µg	35.0	8.5	0.7	36, 43, 26
		5000 µg	47.0	9.5	0.9	56, 48, 37
	DMSO		50.3	2.1		52, 48, 51
	Untreated		49.7	2.1		52, 49, 48
TA 100	Bicyclopyrone SL (A16003E)	3 µg	80.0	3.6	1.0	84, 77, 79
		10 µg	84.7	7.1	1.1	77, 91, 86
		33 µg	89.0	6.0	1.2	83, 95, 89
		100 µg	90.7	8.6	1.2	100, 83, 89
		333 µg	76.7	17.5	1.0	72, 96, 62
		1000 µg	85.0	8.7	1.1	90, 75, 90
		2500 µg	85.7	7.6	1.1	91, 89, 77
		5000 µg	93.0	15.6	1.2	84, 111, 84
	DMSO		76.3	9.0		67, 77, 85
	Untreated		82.3	12.4		68, 90, 89

Study Name: 2129100
 Experiment: 2129100 VV Plate
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 01.10.2020
 Date Counted: 08.10.2020

With metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
WP2 pKM101	Bicyclopyrone SL (A16003E)	3 µg	231.7	11.5	1.1	243, 220, 232
		10 µg	213.3	16.9	1.0	209, 232, 199
		33 µg	182.7	7.5	0.9	183, 190, 175
		100 µg	216.3	5.5	1.0	210, 219, 220
		333 µg	188.7	15.0	0.9	204, 188, 174
		1000 µg	210.7	11.6	1.0	224, 205, 203
		2500 µg	206.3	21.2	1.0	200, 189, 230
		5000 µg	207.0	38.6	1.0	165, 215, 241
		DMSO		207.0	23.5	
	Untreated		234.3	5.9		241, 232, 230
WP2 uvrA pKM101	Bicyclopyrone SL (A16003E)	3 µg	320.3	10.4	1.0	317, 332, 312
		10 µg	298.7	17.9	1.0	303, 314, 279
		33 µg	288.3	11.7	0.9	278, 301, 286
		100 µg	299.0	23.4	1.0	286, 326, 285
		333 µg	311.0	29.5	1.0	343, 305, 285
		1000 µg	284.0	23.3	0.9	305, 259, 288
		2500 µg	302.7	25.1	1.0	324, 309, 275
		5000 µg	301.0	9.5	1.0	296, 312, 295
		DMSO		307.3	4.0	
	Untreated		330.0	11.8		343, 327, 320
TA 1535	2-AA	2.5 µg	199.7	24.2	18.2	174, 203, 222
TA 1537	2-AA	2.5 µg	371.3	37.0	28.6	349, 414, 351
TA 98	2-AA	2.5 µg	2141.0	423.5	42.5	1966, 1833, 2624
TA 100	2-AA	2.5 µg	2437.7	242.6	31.9	2186, 2670, 2457
WP2 pKM101	2-AA	10.0 µg	976.0	31.2	4.7	995, 993, 940
WP2 uvrA pKM101	2-AA	10.0 µg	1567.0	118.4	5.1	1511, 1487, 1703

Key to Positive Controls

2-AA 2-aminoanthracene

TABLE 5 Experiment II: 2129100 HV2 Pre Incubation Without Metabolic Activation

Study Name: 2129100
 Experiment: 2129100 HV2 Pre
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 15.10.2020
 Date Counted: 21.10.2020

Without metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
TA 1535	Bicyclopyrone SL (A16003E)	33 µg	9.7	2.3	1.4	11, 11, 7
		100 µg	6.7	0.6	1.0	7, 7, 6
		333 µg	9.7	2.3	1.4	11, 7, 11
		1000 µg	9.0	2.6	1.3	6, 11, 10
		2500 µg	6.7	2.1	1.0	6, 9, 5
		5000 µg	6.3	2.3	0.9	5, 9, 5
	DMSO Untreated		13.0	3.6		7, 6, 7 10, 17, 12
TA 1537	Bicyclopyrone SL (A16003E)	33 µg	9.7	0.6	0.7	10, 9, 10
		100 µg	12.7	3.8	0.9	10, 11, 17
		333 µg	12.7	2.1	0.9	12, 11, 15
		1000 µg	13.3	2.3	1.0	16, 12, 12
		2500 µg	11.3	0.6	0.8	12, 11, 11
		5000 µg	14.0	2.0	1.0	14, 12, 16
	DMSO Untreated		13.7	2.5		16, 11, 14 16, 15, 10
TA 98	Bicyclopyrone SL (A16003E)	33 µg	22.3	2.3	0.7	21, 25, 21
		100 µg	30.0	7.0	0.9	23, 37, 30
		333 µg	29.3	7.5	0.9	25, 25, 38
		1000 µg	23.7	7.0	0.7	17, 31, 23
		2500 µg	22.3	4.9	0.7	20, 28, 19
		5000 µg	25.7	7.8	0.8	17, 32, 28
	DMSO Untreated		32.7	2.9		31, 31, 36 36, 27, 20
TA 100	Bicyclopyrone SL (A16003E)	33 µg	101.3	11.0	1.1	110, 89, 105
		100 µg	101.7	0.6	1.1	102, 102, 101
		333 µg	82.0	14.4	0.9	98, 70, 78
		1000 µg	86.0	8.0	0.9	78, 86, 94
		2500 µg	81.7	11.9	0.9	78, 72, 95
		5000 µg	76.3	7.1	0.8	84, 75, 70
	DMSO Untreated		93.0	4.4		98, 90, 91 112, 105, 100

Study Name: 2129100
 Experiment: 2129100 HV2 Pre
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 15.10.2020
 Date Counted: 21.10.2020

Without metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
WP2 pKM101	Bicyclopyrone SL (A16003E)	33 µg	268.0	13.2	1.1	273, 253, 278
		100 µg	253.0	8.5	1.0	245, 252, 262
		333 µg	232.0	11.8	0.9	242, 219, 235
		1000 µg	241.0	26.9	1.0	272, 224, 227
		2500 µg	236.3	30.9	1.0	264, 203, 242
		5000 µg	218.3	21.9	0.9	243, 201, 211
	DMSO Untreated		245.7	24.9		258, 217, 262
		286.0	13.9		295, 270, 293	
WP2 uvrA pKM101	Bicyclopyrone SL (A16003E)	33 µg	324.3	7.6	1.0	333, 319, 321
		100 µg	289.0	25.0	0.9	289, 314, 264
		333 µg	311.0	9.5	1.0	320, 312, 301
		1000 µg	315.7	7.6	1.0	307, 321, 319
		2500 µg	298.7	19.0	0.9	317, 279, 300
		5000 µg	265.7	13.6	0.8	264, 280, 253
	DMSO Untreated		325.7	18.6		331, 305, 341
		355.0	26.2		325, 373, 367	
TA 1535	NaN3	10 µg	1047.3	28.9	157.1	1073, 1053, 1016
TA 1537	4-NOPD	50 µg	105.0	12.3	7.7	110, 114, 91
TA 98	4-NOPD	10 µg	702.0	56.4	21.5	738, 731, 637
TA 100	NaN3	10 µg	1578.0	136.5	17.0	1430, 1605, 1699
WP2 pKM101	MMS	2.0 µL	2830.0	23.5	11.5	2841, 2846, 2803
WP2 uvrA pKM101	MMS	2.0 µL	2797.3	80.5	8.6	2712, 2808, 2872

Key to Positive Controls

NaN3	sodium azide
4-NOPD	4-nitro-o-phenylene-diamine
MMS	methyl methane sulfonate

TABLE 6 Experiment II: 2129100 HV2 Pre Incubation With Metabolic Activation

Study Name: 2129100
 Experiment: 2129100 HV2 Pre
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 15.10.2020
 Date Counted: 21.10.2020

With metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
TA 1535	Bicyclopyrone SL (A16003E)	33 µg	12.0	2.6	1.2	10, 11, 15
		100 µg	10.3	1.5	1.0	9, 10, 12
		333 µg	9.3	0.6	0.9	10, 9, 9
		1000 µg	9.0	3.0	0.9	6, 12, 9
		2500 µg	11.3	3.2	1.1	15, 9, 10
		5000 µg	12.3	2.5	1.2	15, 10, 12
	DMSO Untreated		10.0	3.5		6, 12, 12
		10.7	1.5		12, 11, 9	
TA 1537	Bicyclopyrone SL (A16003E)	33 µg	12.7	2.1	0.9	12, 11, 15
		100 µg	12.3	1.5	0.9	14, 12, 11
		333 µg	15.0	0.0	1.1	15, 15, 15
		1000 µg	13.0	2.6	1.0	15, 14, 10
		2500 µg	14.3	4.5	1.1	19, 14, 10
		5000 µg	11.7	0.6	0.9	11, 12, 12
	DMSO Untreated		13.3	2.9		10, 15, 15
		17.3	2.3		16, 16, 20	
TA 98	Bicyclopyrone SL (A16003E)	33 µg	39.7	1.5	1.0	41, 40, 38
		100 µg	38.0	6.2	0.9	31, 40, 43
		333 µg	34.7	11.9	0.9	25, 31, 48
		1000 µg	38.7	6.7	1.0	46, 37, 33
		2500 µg	36.7	9.0	0.9	32, 31, 47
		5000 µg	35.7	11.8	0.9	43, 42, 22
	DMSO Untreated		40.7	3.5		44, 37, 41
		43.0	9.5		48, 32, 49	
TA 100	Bicyclopyrone SL (A16003E)	33 µg	95.0	9.6	1.1	102, 99, 84
		100 µg	92.3	8.3	1.1	83, 95, 99
		333 µg	100.3	7.0	1.2	93, 107, 101
		1000 µg	96.7	9.7	1.1	86, 99, 105
		2500 µg	91.3	10.0	1.1	99, 80, 95
		5000 µg	107.0	6.2	1.2	109, 100, 112
	DMSO Untreated		86.7	4.2		88, 82, 90
		112.0	7.0		117, 104, 115	

Study Name: 2129100
 Experiment: 2129100 HV2 Pre
 Assay Conditions:

Study Code: ICCR 2129100
 Date Plated: 15.10.2020
 Date Counted: 21.10.2020

With metabolic activation

Strain	Compound	Concentration per plate	Mean revertants per plate	Standard Deviation	Ratio treated / solvent	Individual revertant colony counts
WP2 pKM101	Bicyclopyrone SL (A16003E)	33 µg	266.7	13.7	1.0	279, 269, 252
		100 µg	240.7	23.7	0.9	226, 268, 228
		333 µg	242.0	10.5	0.9	231, 243, 252
		1000 µg	259.7	16.2	1.0	270, 241, 268
		2500 µg	249.7	33.7	0.9	225, 236, 288
		5000 µg	266.3	9.7	1.0	258, 277, 264
	DMSO Untreated		271.0	30.3		288, 236, 289
		289.7	2.9		288, 288, 293	
WP2 uvrA pKM101	Bicyclopyrone SL (A16003E)	33 µg	348.3	27.2	1.1	362, 366, 317
		100 µg	332.0	31.6	1.0	299, 335, 362
		333 µg	325.0	17.1	1.0	307, 341, 327
		1000 µg	331.7	21.5	1.0	354, 330, 311
		2500 µg	330.0	18.2	1.0	341, 340, 309
		5000 µg	326.3	15.6	1.0	328, 310, 341
	DMSO Untreated		323.0	8.9		320, 333, 316
		372.0	15.4		389, 368, 359	
TA 1535	2-AA	2.5 µg	259.3	17.6	25.9	279, 245, 254
TA 1537	2-AA	2.5 µg	402.3	43.9	30.2	352, 422, 433
TA 98	2-AA	2.5 µg	3133.7	158.6	77.1	2953, 3250, 3198
TA 100	2-AA	2.5 µg	3501.3	248.2	40.4	3755, 3259, 3490
WP2 pKM101	2-AA	10.0 µg	940.7	11.8	3.5	927, 948, 947
WP2 uvrA pKM101	2-AA	10.0 µg	1859.7	41.8	5.8	1821, 1904, 1854

Key to Positive Controls

2-AA 2-aminoanthracene

APPENDICES SECTION

APPENDIX 1 Historical Control Data

These data represent the laboratory's historical control data from July 2018 until July 2020 representing approx. 600 experiments (WP2 pKM101, WP2 uvrA pKM101 the historical data are based on approx. 80 experiments).

The positive controls that used to compile the historical positive control data correspond to the positive control substances described in Methods; section 3.2.2 (Positive control substances).

Strain		without S9 mix				with S9 mix			
		Mean	SD	Min	Max	Mean	SD	Min	Max
TA 1535	Solvent control	12	2.6	7	22	13	2.5	7	24
	Untreated control	12	2.9	6	26	13	2.8	7	23
	Positive control	1116	141.3	340	1612	346	72.1	170	736
TA1537	Solvent control	11	2.4	6	20	14	2.8	7	28
	Untreated control	11	2.8	5	22	14	3.2	7	30
	Positive control	83	22.1	48	400	286	98.7	82	630
TA 98	Solvent control	28	4.9	13	46	38	6.4	12	62
	Untreated control	29	5.0	14	48	41	6.8	14	64
	Positive control	421	91.2	216	1218	3275	774.9	322	5699
TA 100	Solvent control	127	30.7	63	214	131	30.0	72	214
	Untreated control	135	35.7	64	233	140	34.4	68	217
	Positive control	1759	273.4	511	2588	3566	837.6	553	5444
WP2 pKM 101	Solvent control	248	31.7	171	299	266	33.0	205	315
	Untreated control	269	26.6	212	346	299	28.2	233	345
	Positive control	3343	428.4	2332	4653	1092	257.8	933	2781
WP2uvrA pKM 101	Solvent control	322	31.6	248	388	375	38.5	287	466
	Untreated control	346	28.2	279	403	393	32.6	313	480
	Positive control	3176	468.5	2021	4717	1897	183.2	1270	2464

Mean = mean value of revertants/plate

SD = standard deviation

Min = minimal value

Max = maximal value

APPENDIX 2 Copy of GLP Certificate



Gute Laborpraxis/Good Laboratory Practice

GLP-Bescheinigung/Statement of GLP Compliance (gemäß/according to § 19b Abs. 1 Chemikaliengesetz)



Eine GLP-Inspektion zur Überwachung der Einhaltung der GLP-Grundsätze gemäß Chemikaliengesetz bzw. Richtlinie 2004/9/EG wurde durchgeführt in

Assessment of conformity with GLP according to Chemikaliengesetz and Directive 2004/9/EEC at:

Prüfeinrichtung/Test facility Prüfstandort/Test site

ICCR-Roßdorf GmbH
Institute for Competent Contract Research
In den Leppsteinswiesen 19
64380 Roßdorf

(Unverwechselbare Bezeichnung und Adresse/Unequivocal name and address)

Prüfungen nach Kategorien/Areas of Expertise (gemäß/according ChemVwV-GLP Nr. 5.3/OECD guidance)

- | | |
|-----------------------------------------------------------------------------------------------|----------------------------------------------------|
| 2 Prüfungen zur Bestimmung der toxikologischen Eigenschaften | 2 Toxicity studies |
| 3 Prüfungen zur Bestimmung der erbgutverändernden Eigenschaften (in vitro und in vivo) | 3 Mutagenicity studies |
| 8 Analytische Prüfungen an biologischen Materialien | 8 Analytical and clinical chemistry testing |

22.11.2018, 21.02.2019, 12. bis 14.03.2019
Datum der Inspektion/Date of Inspection
(Tag Monat Jahr/day month year)

Die genannte Prüfeinrichtung befindet sich im nationalen GLP-Überwachungsverfahren und wird regelmäßig auf Einhaltung der GLP-Grundsätze überwacht.

The above mentioned test facility is included in the national GLP Compliance Programme and is inspected on a regular basis.

Auf der Grundlage des Inspektionsberichtes wird hiermit bestätigt, dass in dieser Prüfeinrichtung die oben genannten Prüfungen unter Einhaltung der GLP-Grundsätze durchgeführt werden können.

Based on the inspection report it can be confirmed, that this test facility is able to conduct the aforementioned studies in compliance with the Principles of GLP.

Im Auftrag

Dr. Astrid Brandt, Referentin, Wiesbaden, den **23. Oktober 2019**
(Name und Funktion der verantwortlichen Person/
Name and function of responsible person)



Hessisches Ministerium für Umwelt, Klimaschutz, Landwirtschaft und Verbraucherschutz,
Mainzer Straße 80, D 65189 Wiesbaden
(Name und Adresse der GLP-Überwachungsbehörde/Name and address of the GLP Monitoring Authority)

English name and address of the GLP Monitoring Authority: Hessian Ministry for Environment, Climate Protection, Agriculture and Consumer Protection; Department II 10; P.O. Box 31 09; 65189 Wiesbaden

Translation of seal inscription: Hessian Ministry for Environment, Climate Protection, Agriculture and Consumer Protection

APPENDIX 3 Certificate of S9



CERTIFICATE

ICCR-Roßdorf S9 Preparation Lot No. 030920K

Date of preparation: September 03, 2020

Release date: September 14, 2020


Protein assay: 33 mg protein / ml S9

Sterility: 0 colonies / ml S9 on glucose-minimal-agar


Salmonella typhimurium assay (AMES-test)

Treatment	µl S9 / plate	number of revertants in TA 98
negative	0	28
control	100	33
10 µg/plate	0	69
2-Aminoanthracene	100	2608
10 µg/plate	0	30
Benzo(a)pyrene	100	99

The S9 was obtained from the livers of male Wistar rats which received triple treatments of 80 mg / kg body weight Phenobarbital and β -Naphthoflavone orally on consecutive days. The livers were prepared 24 hours after the last treatment.


Quality Assurance Auditor
ICCR-Roßdorf GmbH

30. SEP. 2020
Date


Dr. Steffen Naumann
Study Director
ICCR-Roßdorf GmbH

30. SEP. 2020
Date

ICCR-Roßdorf GmbH
In den Leppsteinswiesen 19, 64380 Roßdorf, Deutschland
T +49 6154 8070 F +49 6154 83399
Registergericht Darmstadt, HRB 6837, USt-ID DE812333696
Geschäftsführer: Dr. Markus Schulz

SOP Origin TS-SOP S9_21

APPENDIX 4 Certificate of Analysis



Syngenta Crop Protection, LLC
Analytical and Product Chemistry
Greensboro, NC 27409

Certificate of Analysis

A16003E Batch ID 1099852 (ACD9A00004 (ARG))

Test Substance Name:	NOA449280 SL (200)
Common Name:	Bicyclopyrone SL (200)
Material ID:	A16003E
Batch ID:	1099852
Other ID:	ACD9A00004 (ARG)
Source:	DuPont,ARG,Ruta Prov. 33,Km 738,Prov de SantaFe,2170

Chemical Analysis

AI	% w/w	g/L
Bicyclopyrone	18.5	199

Identity of the Active Ingredients: Confirmed

Methodology Used for Characterization: LC, mass spectrometry, oscillating density meter

The Active Ingredient(s) content is within the FAO limits.

Physical Analysis

Property	Value	Units
Density	1.077	g/cm ³

Appearance: Brown Transparent Liquid

Storage Temperature: <30°C

Re-certification Date:End of Aug/2022

If stored under the conditions given above, this test substance can be considered stable until the recertification date is reached.

COA Number: USGR190324

Page 1 of 2

The stability of this test substance will be determined concurrently through reanalysis of material held in inventory under GLP conditions at Syngenta Crop Protection, LLC, Greensboro, NC.


This Certificate of Analysis is summarizing data from a study that has been performed in compliance with Good Laboratory Practices per 40 CFR Part 160. Raw data, documentation, protocols, any amendments to study protocols and reports pertaining to this study are maintained in the Syngenta Crop Protection Archives in Greensboro, NC.

Study Number: USGR190324

Authorization: Gerald Ducatte



Gerald Ducatte
Analytical and Product Chemistry Department



Date

COA Number: USGR190324

Page 2 of 2