

Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram

Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram FS (A23793B) – *In Vitro* Skin Irritation Test in the EPISKIN™ Model

Final Report

TEST GUIDELINE(S):

OECD No. 439 (2021)
EC No 640/2012, B.46. (2012)

AUTHOR(S):

Balázs Orovecz, B.Sc.

COMPLETION DATE:

30 November 2021

PERFORMING LABORATORY: Charles River Laboratories Hungary Kft.
H-8200 Veszprém, Szabadságpuszta, hrsz. 028/1.,
Hungary

LABORATORY PROJECT ID:

Report Number: 21/245-043B
Study Number: 21/245-043B
Task Number: TK0518490

SPONSOR(S):

Syngenta Ltd.
Jealott's Hill International Research Centre
Bracknell, Berkshire, RG42 6EY, United Kingdom

SEGREDO INDUSTRIAL

Este documento é considerado de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo-se em SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

STATEMENT OF DATA CONFIDENTIALITY CLAIMS

The Following Statement Applies To The United States of America:

STATEMENT OF NO DATA CONFIDENTIALITY CLAIMS UNDER SPECIFIED FIFRA PROVISIONS

No claim of confidentiality, on any basis whatsoever, is made for any information contained in this document. I acknowledge that information not designated as within the scope of FIFRA sec. 10(d)(1)(A), (B), or (C) and which pertains to a registered or previously registered pesticide is not entitled to confidential treatment and may be released to the public, subject to the provisions regarding disclosure to multinational entities under FIFRA 10(g).

Company: Syngenta Crop Protection, LLC
410 Swing Road
Post Office Box 18300
Greensboro, NC 27419-8300 USA

Submitter: _____ Date: _____

Syngenta is the owner of this information and data. Syngenta has submitted this material to the United States Environmental Protection Agency specifically under the provisions contained in FIFRA as amended and, hereby, consents to use and disclosure of this material by EPA according to FIFRA. In submitting this material to EPA according to method and format requirements contained in PR Notice 2011-3, we do not waive any protection or right involving this material that would have been claimed by the company if this material had not been submitted to the EPA, nor do we waive any protection or right provided under FIFRA Section 3 (concerning data exclusivity and data compensation) or FIFRA Section 10(g) (prohibiting disclosure to foreign and multinational pesticide companies or their agents).

SEGREDO INDUSTRIAL

Este documento é considerado de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo SEGREDO DE NEGÓCIOS E SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

GOOD LABORATORY PRACTICE COMPLIANCE STATEMENT

This study has been performed in accordance with the study plan and the Principles of Good Laboratory Practice (Hungarian GLP Regulations: 42/2014. (VIII. 19.) EMMI decree of the Ministry of Human Capacities which corresponds to the OECD GLP, ENV/MC/CHEM (98) 17).

I, the undersigned, declare that this report constitutes a true record of the actions undertaken and the results obtained in the course of this study. By virtue of my dated signature I accept the responsibility for the validity of the data.

Signature:



Balázs Orovecz, B.Sc.
Study Director

Date:

30 November 2021

Performing Laboratory:

Charles River Laboratories Hungary Kft.
H-8200 Veszprém, Szabadságpuszta, hrsz. 028/1.,
Hungary

To be completed for USA EPA submission only:

Representative of Submitter/Sponsor:

Date

Submitter/Sponsor: Syngenta Crop Protection, LLC
410 Swing Road
Post Office Box 18300
Greensboro, NC 27419-8300 USA

SEGREDO INDUSTRIAL

Estas informações são confidenciais e de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo
Report Number: 21/245-043B SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei
9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 3 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles
descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

FLAGGING STATEMENT

This page is intentionally left blank. It will be replaced by an appropriate Flagging statement by the sponsor.



SEGREDO INDUSTRIAL

Este documento é considerado de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

QUALITY ASSURANCE STATEMENT

Study Number: 21/245-043B


Study Title: Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram FS (A23793B)
– *In Vitro* Skin Irritation Test in the EPISKIN™ Model

Test Item: Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram FS
(A23793B)

This study has been inspected, and this report audited by the Quality Assurance Unit in compliance with the Principles of Good Laboratory Practice. As far as it can be reasonably established the methods described and the results incorporated in this report accurately reflect the raw data produced during this study.

All inspections, data reviews and the report audit were reported in written form to the study director and to management. The dates of such inspections and of the report audit are given below:

Date of Inspection	Phase(s) Inspected/Audited	Date of report to	
		Management	Study Director
14 October 2021	Study Plan	14 October 2021	14 October 2021
20 October 2021	Treatment	20 October 2021	20 October 2021
05 November 2021	Draft Report	05 November 2021	05 November 2021
19 November 2021	Amendment 1 to the Study Plan	19 November 2021	19 November 2021
29 November 2021	Final Report	29 November 2021	29 November 2021

Signature: 
Eszter Sebestyén, B.Sc.
On Behalf of QA

Date: 30 November 2021

Report Number: 21/245-043B
ESTAS INFORMAÇÕES SÃO CONFIDENCIAIS E DE PROPRIEDADE DA SYNGENTA PROTEÇÃO DE CULTIVOS LTDA., CONSTITUINDO
9.279/96 E DO PARÁGRAFO 2º DO ARTIGO 9º DA LEI 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles
descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

MANAGEMENT STATEMENT

According to the conditions of the research and development agreement between Syngenta Ltd. (as Sponsor) and Charles River Laboratories Hungary Kft. (as Test Facility), the study titled “Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram FS (A23793B) – *In Vitro* Skin Irritation Test in the EPISKIN™ Model” was performed, in compliance with the Principles of Good Laboratory Practice.

Signature: 

Balázs Tóth, Ph.D.
General Manager

Date: 30 November 2021

SEGREDO INDUSTRIAL

Estas informações são confidenciais e de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo Report Number: 21/245-043B DO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 6 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

GENERAL INFORMATION

Contributors

The following contributed to this report in the capacities indicated:

Name	Function
Balázs Orovecz, B.Sc.	Study Director
Balázs Tóth, Ph.D.	General Manager
David J. Esdaile, M.Sc.	Director of Science and Regulatory Affairs
Kata Tóth-Gönczöl, B.Sc.	Assistant Scientist
Eszter Sebestyén, B.Sc.	Quality Assurance Auditor
Carolina Vaccari, M.Sc.	Syngenta Study Manager

Study dates

Study Plan:	15 October 2021
Amendment 1 to the Study Plan:	22 November 2021
Experimental Starting Date:	19 October 2021
Experimental Completion Date:	22 October 2021
Date of Draft Report (non-QA audited):	22 October 2021
Date of Draft Report (QA audited):	08 November 2021
Date of Final Report:	30 November 2021

Note: MTT-pre check was performed on 19 October 2021

Deviations from the Study Plan

There were no deviations from the Study Plan.

Performing laboratory test substance reference number

210563

Other

The study documents and samples:

- study plan and amendment,
- all raw data,
- sample of the test item,
- original study report and any amendments,
- correspondence

will be archived according to the Hungarian GLP regulations and to applicable SOP's in the Archives of Charles River Laboratories Hungary Kft. (H-8200 Veszprém, Szabadságpuszta, hrsz. 028/1., Hungary). This is for a period of 15 years.

After the retention time of 15 years has elapsed all the archived materials listed above will be returned to the Sponsor or retained for a further period if agreed by a contract. Otherwise the materials will be discarded.

SEGREDOES INDUSTRIALIS

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo SEGREDO DE NEGÓCIOS E SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N.º 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 7 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

TABLE OF CONTENTS

STATEMENT OF DATA CONFIDENTIALITY CLAIMS	2
GOOD LABORATORY PRACTICE COMPLIANCE STATEMENT	3
FLAGGING STATEMENT	4
QUALITY ASSURANCE STATEMENT	5
MANAGEMENT STATEMENT	6
GENERAL INFORMATION	7
TABLE OF CONTENTS	8
1.0 EXECUTIVE SUMMARY	11
1.1 Study Design	11
1.2 Results	11
1.3 Conclusion.....	11
2.0 INTRODUCTION	12
2.1 Purpose.....	12
2.2 Guidelines	12
2.3 Test Facility.....	12
3.0 MATERIALS AND METHODS	13
3.1 Test Substance.....	13
3.2 Identification and Receipt	13
3.3 Test Item Preparation	13
3.4 Subsidiary Materials.....	13
3.4.1 Negative control	14
3.4.2 Positive control.....	14
3.5 Test System	14
3.5.1 Human skin	14
3.5.2 Quality control	15
3.5.3 Justification for selection of the test system	15
3.5.4 Kit contents	15
3.5.5 Number of replicate wells	15
3.5.6 Kit reception.....	15
3.5.7 Storage.....	16
3.6 Additional Materials.....	16
3.6.1 MTT solution	16
3.6.2 Acidified isopropanol.....	16

SEGREDO INDUSTRIAL

Este documento contém informações de propriedade da Syngenta Proteção de Cultivos Ltda., constituindo SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 8 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.6.3	Chemicals used in the experiment.....	16
3.7	Indicator for Potential False Viability.....	16
3.7.1	Check-method for possible direct MTT reduction with test item	17
3.7.2	Check-method to detect the colouring potential of test item	17
3.7.3	Additional control tissues used in respect of colour interference potential (NSC _{living})	17
3.8	Performance of the Study.....	17
3.8.1	Pre-incubation (Day [-1]).....	18
3.8.2	Application and rinsing (Day 0).....	18
3.8.3	MTT test (Day 2)	19
3.8.4	Formazan extraction (Day 2)	19
3.8.5	Cell viability measurements (Day 2).....	19
3.9	Calculations of Viability Percentages	19
3.9.1	Data calculation for normal test items	19
3.9.2	Data calculation for test items having MTT-interacting potential	20
3.9.3	Data calculation for test items having colouring potential.....	21
3.9.4	Data calculation for substance having both MTT interacting potential and colouring potential	22
3.10	Validity of the Test.....	22
3.11	Interpretation of Test Results	22
4.0	RESULTS AND DISCUSSION	23
4.1	Additional Controls	23
4.2	Viability Results.....	23
4.3	Validity of the Test.....	24
4.4	Discussion	24
5.0	CONCLUSIONS	24
6.0	REFERENCES	24
7.0	DISTRIBUTION OF THE FINAL REPORT	25
TABLES SECTION		26
TABLE 1	Optical Density (OD) and the Calculated Non Specific Colour % (NSC _{living} %) of the Additional Control Tissues	27
TABLE 2	Optical Density (OD) and the Calculated Relative Viability % of the Samples	28
APPENDICES SECTION		29
APPENDIX 1	Certificate of Analysis.....	30
APPENDIX 2	Copy of the Test Kit Quality Control.....	32
APPENDIX 3	Historical Control Data	33



SEGREDO INDUSTRIAL

Este documento é considerado Segredo de propriedade da Syngenta Proteção de Cultivos Ltda., constituidos do SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 10 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

1.0 EXECUTIVE SUMMARY

1.1 Study Design

An *in vitro* skin irritation test was conducted on difenoconazole/fludioxonil/metalaxyl-m/cyclobutifluram FS (A23793B) in a reconstructed human epidermis model. The EpiSkin™ model is designed to predict and classify the irritation potential of chemicals by measuring its cytotoxic effect as reflected in the MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, Thiazolyl blue; CAS number: 298-93-1) assay. The irritation potential of the test item was evaluated according to the OECD guideline No. 439 [1].

Disks of EpiSkin™ (three units) were treated with the test item and incubated for 15 minutes at room temperature. Exposure of the test item to the EpiSkin™ surface was terminated by rinsing the units with Phosphate Buffered Saline (PBS). The epidermis units were then incubated at 37°C for 42 hours in an incubator with 5% CO₂ in a >95% relative humidified (RH) atmosphere. After the 42-hour incubation, MTT solution was added to the units and incubated for a further 3 hours to determine cell viability. The precipitated formazan crystals were then extracted using acidified isopropanol and quantified spectrophotometrically at 570 nm.

The negative control epidermis units were treated with PBS, whilst the positive control epidermis units were treated with 5% (w/v) sodium dodecyl sulphate (SDS) (three units / control). Two additional disks were used to provide an estimate of colour contribution (NSC_{living}) from the test item. For each treated tissue, the viability was expressed as a percentage relative to the negative control. If the mean relative cell viability after 15 minutes of exposure and 42 hours post incubation is less or equal (≤) to 50% of the negative control, the test item is considered to be an irritant to skin.

1.2 Results

Following exposure to difenoconazole/fludioxonil/metalaxyl-m/cyclobutifluram FS (A23793B), the mean cell viability was 123.9% compared to the negative control. This is above the threshold of 50%, therefore the test item was considered as being non-irritant to skin under the conditions of this assay. The experiment met the validity criteria, and therefore the study was considered to be valid.

1.3 Conclusion

In conclusion, under the conditions of this *in vitro* EpiSkin™ irritation assay, the results indicate that difenoconazole/fludioxonil/metalaxyl-m/cyclobutifluram FS (A23793B) is non-irritant to skin.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

2.0 INTRODUCTION

2.1 Purpose

The skin irritation potential of a test item may be predicted by measurement of its cytotoxic effect, as reflected in the MTT assay, on the EpiSkin™ reconstituted human epidermis [1-3]. This method is approved by international regulatory agencies as a replacement for the identification of irritants in the *in vivo* rabbit skin assay (OECD No. 404) [4].

The test is designed to predict and classify the skin irritant potential of chemicals/formulations/products/mixtures according to chemical safety regulations, using the reconstructed human epidermis model EpiSkin™ and parameters related to skin irritation.

EpiSkin™ is a three-dimensional human skin model comprised of a reconstructed epidermis with a functional stratum corneum. Its use for skin irritation testing involves topical application of test materials to the surface of the epidermis, and the subsequent assessment of their effects on cell viability. Cell viability is determined based on cellular mitochondrial dehydrogenase activity, measured by MTT reduction and conversion into a blue formazan salt that is quantitatively measured after extraction from tissues (Faller C. et al., 2002, Mosmann T., 1983) [5, 6]. The reduction in cell viability in treated tissues is compared to negative controls and expressed as a percentage. The percent reduction in cell viability is used to predict the irritation potential of the test item.

The OECD No. 439 is specifically a test for dermal irritation; it is not a test for corrosivity. A negative or positive result in this assay does not necessarily mean that the test item is non-corrosive.

2.2 Guidelines

The study followed the procedures indicated by the following internationally accepted guidelines and recommendations:

- OECD Guidelines for Testing of Chemicals, Section 4, No. 439, “*In Vitro* Skin Irritation: Reconstructed Human *Epidermis* Test Method”, 14 June 2021
- Commission Regulation (EC) No 640/2012 of 6th July 2012 amending, for the purpose of its adaptation to technical progress, Regulation (EC) No 440/2008, B.46.

2.3 Test Facility

This study was performed in an AAALAC-accredited laboratory. The Institutional Animal Care and Use Committee (IACUC) of Charles River Laboratories Hungary Kft. monitored the conduct of the study.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.0 MATERIALS AND METHODS

3.1 Test Substance

Information supplied by the Sponsor:

Name: Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram FS (A23793B)

Batch number: 1200767

Design code: A23793B

Active ingredient content*: Difenoconazole 5.45% w/w 64.0 g/L, fludioxonil 4.37% w/w 51.3 g/L, metalaxyl-M 4.31% w/w 50.6 g/L, cyclobutifluram 21.0% w/w 247 g/L

Appearance: Red liquid

Recertification date: 31 August 2024

Storage conditions: Room temperature (15-25°C, ≤70% relative humidity)

Safety precautions: Routine safety precautions (gloves, goggles, face mask, lab coat) for unknown materials were applied to ensure personnel health and safety.

* No adjustment for active ingredient content was applied as agreed by the Sponsor.

A Certificate of Analysis supplied by the Sponsor is given in Appendix 1.

The integrity of supplied data relating to the identity, purity and stability of the test material was the responsibility of the Sponsor.

3.2 Identification and Receipt

The test item of a suitable active ingredient content together with all precautions required in the handling and disposal of the test item were supplied by the Sponsor. The identification of the test item was made on the basis of the information provided by Sponsor in the Pharmacy of Charles River Laboratories Hungary Kft.

3.3 Test Item Preparation

The test item was applied in its original form (undiluted).

3.4 Subsidiary Materials

Positive and negative controls were included in the experiment. Furthermore, as the test item was coloured, two additional control tissue samples were used in the experiment for determination of the non-specific colour (as detailed in 3.7.2.).

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.4.1 Negative control

Phosphate Buffered Saline:

Abbreviation: PBS
Supplier: Life Technologies Corporation
Batch number: 2215013
Expiry date: 31 December 2022

3.4.2 Positive control

5% (w/v) Sodium Dodecyl Sulphate solution (SDS):

SDS (5% w/v aqueous solution) was freshly prepared in the testing laboratory.

The following chemicals were used for the preparation of the positive control solution:

Sodium Dodecyl Sulphate:

Supplier: Sigma-Aldrich
Batch number: STBJ9692
Expiry date: 31 October 2026

Distilled water:

Supplier: B. Braun Pharmaceuticals SA
Batch number: 11511Y25-2
Expiry date: 31 March 2024

3.5 Test System

3.5.1 Human skin

EpiSkin™ (Manufacturer: SkinEthic, France, Batch No.: 21-EKIN-042, Expiry Date: 25 October 2021) is a three-dimensional human epidermis model. Adult human-derived epidermal keratinocytes are seeded on a dermal substitute consisting of a collagen type I matrix coated with type IV collagen. A highly differentiated and stratified epidermis model is obtained after 13-day culture period comprising the main basal, supra basal, spinous and granular layers and a functional stratum corneum (Tinois et al., 1994) [7]. Its use for skin irritation testing involves topical application of test materials to the surface of the epidermis, and the subsequent assessment of their effects on cell viability.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui-se de SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei nº 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.5.2 Quality control

EpiSkin™ kits are manufactured according to defined quality assurance procedures (certified ISO 9001). All biological components of the epidermis and the kit culture medium have been tested for the presence of viruses, bacteria and mycoplasma.

The quality of the final product is assessed by undertaking a MTT cell viability test and a cytotoxicity test with sodium dodecyl sulphate (SDS). These quality control experiments were conducted at SkinEthic laboratories (supplier of the EpiSkin™ test kits used in the present study) and are documented in Appendix 2.

3.5.3 Justification for selection of the test system

The EpiSkin™ has been validated for irritation testing in an international validation study [3] and its use is recommended by the relevant OECD guideline for irritation testing (OECD No. 439); therefore, it was considered to be suitable for this study.

3.5.4 Kit contents

Units: EpiSkin™ plate containing up to 12 reconstructed epidermis units (area: 0.38 cm²) each reconstructed epidermis is attached to the base of a tissue culture vessel with an O-ring set and maintained on nutritive agar for transport.

Plate: 12-well assay plate

Punch: EpiSkin™ biopsy punch for easy sampling of epidermis

Medium: A flask of sterile "Maintenance Medium"
(Batch No.: 21-MAIN3-038; Exp. Date: 27 October 2021)
A flask of sterile "Assay Medium"
(Batch No.: 21-ESSC-037; Exp. Date: 27 October 2021)

3.5.5 Number of replicate wells

In this assay, three replicates were used for the test item. Three negative controls and three positive controls were also run in the assay. As the test item was coloured, two additional test item-treated living tissues were used for the non-specific optical density (OD) evaluation.

3.5.6 Kit reception

The pH of the agar medium used for transport was checked by checking the colour of the medium:

- orange colour = good
- yellow or violet colour = not acceptable

The colour of the temperature indicator was inspected to verify that the kit has not been exposed to a temperature above 40°C (the colour change is irreversible, independent of the length of the period above 40 °C):

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

- white colour = good
- grey or black colour = not acceptable

3.5.7 Storage

The EpiSkin™ kits were kept in their packaging at 37°C, the Assay Medium and Maintenance Medium supplied with the kits were stored at 2-8°C until the initiation of the test.

3.6 Additional Materials

3.6.1 MTT solution

MTT [3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, Thiazolyl blue; CAS number 298-93-1] was diluted in phosphate buffered saline (PBS) at a final concentration of 3 mg/mL (MTT stock solution). The obtained stock solution (prepared on 19 October 2021) was stored in a refrigerator (2-8°C) protected from light. It was diluted with pre-warmed (37°C) Assay Medium to a final concentration of 0.3 mg/mL (MTT working solution) immediately before use.

3.6.2 Acidified isopropanol

Isopropanol was acidified with HCl acid to achieve a final concentration of 0.04M HCl (1.8 mL of 12M HCl acid was diluted in 500 mL isopropanol, or similar ratio was applied). The solution was prepared on the day of use.

3.6.3 Chemicals used in the experiment

The chemicals used in this experiment are summarised in the following table:

Chemical	Supplier	Batch Number	Expiry date
MTT	VWR International Ltd.	19J656213	31 December 2021
Isopropanol (2-Propanol)	VWR International Ltd.	19D244010	23 April 2024
Hydrochloric acid (37% HCl)	VWR International Ltd.	18J094018	03 October 2023
Phosphate buffered saline (PBS)	Life Technologies Corporation	2215013	31 December 2022

3.7 Indicator for Potential False Viability

Optical properties of the test material or its chemical action on MTT may interfere with the assay leading to a false estimate of cell viability. This may occur when the test item is not completely removed from the tissue by rinsing or when it penetrates the epidermis during the exposure period. If the test material directly acts on MTT (MTT-reducer), is naturally coloured, or becomes coloured during tissue treatment, additional controls should be used to

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 16 of 34

Todos os infratores poderão ser processados civil e criminalmente

detect and correct for test item interference with the cell viability measurement. Methods indicating how to correct direct MTT reduction and interferences by colouring agents are detailed in the following paragraphs.

3.7.1 Check-method for possible direct MTT reduction with test item

10 µL of test item was added to 2 mL MTT working solution and mixed. The mixture was incubated at 37°C for 3 hours protected from light, and then any colour change was recorded:

-Test items which directly react with MTT:	blue or purple
-Test items which do not react with MTT:	another colour

After the test item difenoconazole/fludioxonil/metalaxyl-m/cyclobutifluram FS (A23793B) had been incubated with MTT for three hours, red colour of the mixture was detected in the test tube. Thus, the test item did not react with MTT and therefore, additional controls were not used in the experiment.

3.7.2 Check-method to detect the colouring potential of test item

Prior to treatment, the test item was evaluated for its intrinsic colour or ability to become coloured in contact with water and/or isopropanol** (simulating a tissue humid environment). As the test item had an intrinsic colour, the use of additional control tissues (determination of the non-specific colour percentage) was necessary to evaluate the ability of the test item to stain the epidermis.

**Note: Water is the environment during exposure, isopropanol is the extracting solution.

3.7.3 Additional control tissues used in respect of colour interference potential (NSC_{living})

In addition to the normal procedure (3.9.1), two additional test item-treated living tissues were used for the non-specific OD evaluation. These tissues followed the same test item application and all steps as the other tissues, except for the MTT step: MTT incubation was replaced by incubation with fresh Assay Medium to mimic the amount of colour from the test item that may be present in the test disks. OD reading was conducted following the same conditions as for the other tissues.

3.8 Performance of the Study

Procedures described in sections 3.8.1., 3.8.2. and 3.8.3. were performed under aseptic conditions (in sterile hood using sterile equipment).

SEGREDO INDUSTRIAL

Report Number: 21/245-043B
Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui
SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei
9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 17 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles
descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.8.1 Pre-incubation (Day [-1])

The Maintenance Medium was pre-warmed to 37°C. The appropriate number of wells in an assay plate were filled with the pre-warmed medium (2 mL per well). The epidermis units were placed with the media below them, in contact with the epidermis into each prepared well and then incubated overnight (at least 18 hours) at 37°C in an incubator with 5 % CO₂, in a >95 RH % humidified atmosphere.

3.8.2 Application and rinsing (Day 0)

Test Item

10 µL of test item was applied evenly to each of three test units and each additional control skin unit. The 10 µL test item was sufficient to cover the epidermal surface.

Negative and positive controls

10 µL of negative control (PBS) or positive control (5% (w/v) SDS solution) was added to each skin unit by using a suitable pipette. Chemicals were spread gently with the pipette tip in order to cover evenly all the epidermal surface if necessary (without damaging the epidermis).

Note: The negative and positive controls were also part of concurrent study (Charles River Laboratories Hungary Kft. studies codes: 21/200-043B and 21/235-051B) performed in the same experimental period using the same batch of chemicals and same batch of skin units.

Incubation with Test Item

The plates with the treated epidermis units were incubated for the exposure time of 15 minutes (± 0.5 minute) at room temperature (22.9-24.3°C).

Rinsing

After the 15 minutes incubation time, the EpiSkin™ units were removed and rinsed thoroughly with PBS (2*25mL) to remove as much of the remaining test material as possible from the epidermal surface. The remaining PBS was removed from the epidermal surface with a pipette (without touching the epidermis).

Note: test item was stuck on epidermal surfaces.

After rinsing the units were placed into the plate wells with fresh pre-warmed Maintenance Medium (2 mL/well) and then incubated for 42 hours (±1 hour) at 37°C in an incubator with 5% CO₂, in a >95 RH % humidified atmosphere.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.8.3 MTT test (Day 2)

After the 42 hours incubation, all EpiSkin™ units (except the two colour control units, which were incubated with Assay Medium) were transferred into the MTT working solution filled wells (2 mL of 0.3 mg/mL MTT per well). Then, all transferred EpiSkin™ units were incubated for 3 hours at 37°C in an incubator with 5% CO₂ protected from light, in a >95 RH % humidified atmosphere.

3.8.4 Formazan extraction (Day 2)

After the incubation with MTT, a formazan extraction was undertaken. A uniform-size disk of epidermis was cut from each skin unit (this involved the maximum area of the disk) using a biopsy punch (supplied as part of the kit). The epidermis was separated with the aid of forceps and both parts (epidermis and collagen matrix) were placed into a tube containing 500 µL acidified isopropanol (one tube corresponded to one well of the assay plate).

The capped tubes were thoroughly mixed by using a vortex mixer to achieve a good contact of all of the material and the acidified isopropanol, and then incubated for about two hours at room temperature protected from light with gentle agitation (~150 rpm) for formazan extraction.

A blank sample consisting of an empty tube filled with 2 mL of acidified isopropanol was processed in parallel.

3.8.5 Cell viability measurements (Day 2)

Following the formazan extraction, 2×200 µL samples from each tube were placed into the wells of a 96-well plate (labelled appropriately). The OD (optical density or absorbance) of the samples was measured using a plate reader at 570 nm. The mean of 6 wells of acidified isopropanol solution (200 µL/well) was used as blank.

The proper status of the instrument was verified by measuring a Verification plate (Manufacturer: Thermo Fisher Scientific, Catalogue Number: 240 72800, Serial Number: 0920-14, Date of calibration: 14 October 2020, calibration is valid until October 2022) at the required wavelength on each day before use.

3.9 Calculations of Viability Percentages

3.9.1 Data calculation for normal test items

Blank:

- The mean of the six blank OD values was calculated

Negative control:

- Individual negative control OD values (NC_{raw}) were corrected with the mean blank OD:

SEGREDOS INDUSTRIALIS

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

OD Negative Control (OD_{NC}) = OD_{NCraw} – OD_{blank mean}

- The mean corrected OD of the 3 negative control samples was calculated: this value corresponds to 100% viability

Positive control:

- Individual positive control OD values (PC_{raw}) were corrected with the mean blank OD:

OD Positive Control (OD_{PC}) = OD_{PCraw} – OD_{blank mean}

- The mean corrected OD of the 3 positive control samples was calculated
- The viability % for each positive control replicate was calculated relative to the mean negative control:

Positive Control 1 % = (OD_{PC1} / mean OD_{NC}) × 100

Positive Control 2 % = (OD_{PC2} / mean OD_{NC}) × 100

Positive Control 3 % = (OD_{PC3} / mean OD_{NC}) × 100

- The mean value of the 3 individual relative viability % for positive control was calculated:

Mean PC % = (PC1 % + PC2 % + PC3 %) / 3

Test item:

- Individual test item OD values (TT_{raw}) were corrected with the mean blank OD:

OD Treated Tissue (OD_{TT}) = OD_{TTraw} – OD_{blank mean}

- The mean corrected OD of the 3 test item samples was calculated
- The viability % for each test item replicate was calculated relative to the mean negative control:

Treated Tissue 1 % = (OD_{TT1} / mean OD_{NC}) × 100

Treated Tissue 2 % = (OD_{TT2} / mean OD_{NC}) × 100

Treated Tissue 3 % = (OD_{TT3} / mean OD_{NC}) × 100

- The mean value of the 3 individual relative viability % for test item was calculated:

Mean TT % = (TT1 % + TT2 % + TT3 %) / 3

3.9.2 Data calculation for test items having MTT-interacting potential

Test items that interfere with MTT can produce non-specific reduction of the MTT. In this case, additional control samples are used to determine the OD value derived from non-specific reduction of the MTT. The measured OD value is corrected by the result of the additional controls before calculation of viability % as detailed below:

- Non-specific MTT reduction calculation (NSMTT%):

NSMTT (%) = [(OD_{KT} - OD_{KNC}) / OD_{NC}] × 100

OD_{KNC}: negative control killed tissues OD

OD_{KT}: test item treated killed tissues OD

OD_{NC}: negative control OD

If NSMTT% is $\leq 30\%$, then true MTT metabolic conversion (TOD_{TT}) is undertaken as follows:

TOD_{TT} = [OD_{TT} - (OD_{KT} - OD_{KNC})]

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N.º 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

OD_{TT}: test item treated viable tissues

- The relative viability (RV%) for each test item replicate is calculated relative to the mean negative control:

$$RV\ 1\ \% = [TOD_{TT1} / \text{mean } OD_{NC}] \times 100$$

$$RV\ 2\ \% = [TOD_{TT2} / \text{mean } OD_{NC}] \times 100$$

$$RV\ 3\ \% = [TOD_{TT3} / \text{mean } OD_{NC}] \times 100$$

- The mean value of the 3 individual relative viability % for test item is calculated:

$$\text{Mean Relative Viability \%} = (RV\ 1\ \% + RV\ 2\ \% + RV\ 3\ \%) / 3$$

If NSMTT% is > 30% relative to the negative control, then additional steps must be undertaken if possible, or the test item must be considered as incompatible with the test.

3.9.3 Data calculation for test items having colouring potential

For test items detected as able to stain the tissues the non-specific OD due to the residual chemical colour (unrelated to mitochondrial activity) is evaluated and subtracted before calculation of the “true” viability % as detailed below:

- Non-Specific Colour % with viable tissues (NSC_{living} %):

$$NSC_{living}\ \% = (\text{mean } OD_{CTV} / \text{mean } OD_{NC}) \times 100$$

OD_{CTV}: test substance treated viable tissues (not incubated with MTT)

OD_{NC}: negative control OD (incubated with MTT)

If NSC_{living} % is \leq 5 % then the normal calculation mode is used (see 3.9.1).

If NSC_{living} % is > 5% and \leq 30%, then true MTT metabolic conversion (TOD_{TT}) is undertaken as follows.

$$TOD_{TT} = [OD_{TT} - \text{mean } OD_{CTV}]$$

OD_{TT}: test substance treated viable tissue (incubated with MTT)

OD_{CTV}: test substance treated viable tissue (not incubated with MTT)

- The relative viability (RV%) for each test item replicate is calculated relative to the mean negative control:

$$RV\ 1\ \% = [TOD_{TT1} / \text{mean } OD_{NC}] \times 100$$

$$RV\ 2\ \% = [TOD_{TT2} / \text{mean } OD_{NC}] \times 100$$

$$RV\ 3\ \% = [TOD_{TT3} / \text{mean } OD_{NC}] \times 100$$

- The mean value of the 3 individual relative viability % for test item is calculated:

$$\text{Mean Relative Viability \%} = (RV\ 1\ \% + RV\ 2\ \% + RV\ 3\ \%) / 3$$

If NSC_{living} % is > 30 % relative to the negative control, additional steps must be undertaken if possible, or the test substance must be considered as incompatible with the test.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui-se em SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

3.9.4 Data calculation for substance having both MTT interacting potential and colouring potential

For test substances detected as able to stain the tissues (3.9.3) and interfere with MTT (3.9.2), a third set of controls is also required before calculation of the “true” viability %.

- Non-Specific Colour % with killed tissues (NSC_{killed} %):

$$NSC_{killed} \% = (mean\ OD_{CTK} / mean\ OD_{NC}) \times 100$$

OD_{CTK}: test substance treated killed tissues (not incubated with MTT)

OD_{NC}: negative control OD (incubated with MTT)

In that case the true MTT metabolic conversion (TOD_{TT}) is undertaken as follows:

$$TOD_{TT} = [OD_{TT} - (OD_{KT} - OD_{KNC}) - mean\ OD_{CTV} + mean\ OD_{CTK}]$$

OD_{TT}: test substance treated viable tissues (incubated with MTT)

OD_{KT}: test substance treated killed tissues OD

OD_{KNC}: negative control killed tissues OD

OD_{CTV}: test substance treated viable tissues (not incubated with MTT)

OD_{CTK}: test substance treated killed tissues (not incubated with MTT)

- The % relative viability (% RV) for each test substance replicate is calculated relative to the mean negative control:

$$RV\ 1\ \% = [TOD_{TT1} / mean\ OD_{NC}] \times 100$$

$$RV\ 2\ \% = [TOD_{TT2} / mean\ OD_{NC}] \times 100$$

$$RV\ 3\ \% = [TOD_{TT3} / mean\ OD_{NC}] \times 100$$

- The mean value of the 3 individual relative viability % for test substance is calculated:

$$Mean\ Relative\ Viability\ \% = (RV\ 1\ \% + RV\ 2\ \% + RV\ 3\ \%) / 3$$

3.10 Validity of the Test

The mean OD value of the three negative control tissues should be between 0.6 and 1.5 and the standard deviation value (SD) of the % viability values should be ≤ 18 .

The acceptable mean percentage viability range for the three positive controls is 0-40% and the standard deviation value (SD) of the % viability values should be ≤ 18 .

The SD calculated from individual % tissue viability values of the three test item treated replicates should be ≤ 18 .

The mean OD value of the blank samples (acidified isopropanol) should be <0.1 .

3.11 Interpretation of Test Results

The irritation potential of test items can be classified according to the United Nations Globally Harmonized System of Classification and Labelling of Chemicals [9], and a similar system is used in CLP [11,12]. In the present study, the irritancy potential of test items is

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui-se em SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

predicted by the mean tissue viability of tissues exposed to the test item. The test chemical is identified as requiring classification and labelling according to UN GHS (Category 2 or Category 1) if the mean relative viability of three individual tissues after 15 minutes of exposure to the test item and 42 hours post incubation is less or equal (\leq) to 50% of the mean viability of the negative controls. In case the test chemical is found to be non-corrosive and shows tissue viability after exposure and post-treatment incubation less than or equal (\leq) to 50%, the test chemical is considered to be irritant to skin in accordance with UN GHS Category 2. The test item may be considered to be non-irritant to skin in accordance with UN GHS (No Category), if the mean relative viability of three individual tissues after 15 minutes of exposure to the test item and 42 hours post incubation is more than ($>$) 50% of the mean viability of the negative controls.

The prediction model (PM) is described below:

Criteria for <i>In Vitro</i> interpretation	Classification
	UN GHS
Mean tissue viability % is \leq 50 %	Category 2 or Category 1
Mean tissue viability % is $>$ 50 %	Non-Irritant*

*Note: If there is clear evidence that the test item is not corrosive, then it can be determined as No Category according to the UN GHS. It is plausible that some weaker corrosives could be classified as non-irritant in this *in vitro* assay.

4.0 RESULTS AND DISCUSSION

4.1 Additional Controls

Red colour was observed after three hours of incubation of the test item in MTT working solution, thus the test material did not interact with MTT. Therefore, additional controls and data calculations were not necessary. The false estimation of viability can be excluded.

As the test item was coloured, two additional test item-treated living tissues were used for the non-specific OD evaluation. The mean optical density (measured at 570 nm) of tissues was 0.013, non-specific colour percent was calculated as 1.7% (see Table 1). This value was below 5%, therefore an additional data calculation to account for non-specific colouring was not necessary.

4.2 Viability Results

The results of the optical density (OD) measured at 570 nm for the test item, positive and negative controls and the calculated relative viability percentage values (RV%) are presented in Table 2. The OD values were used to calculate the mean relative viability of the test item treated skin samples, which was shown to be 123.9%.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N.º 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 23 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

4.3 Validity of the Test

After receipt, the two indicators included in the package of the delivered kits (reflecting the storage temperature history and the pH) were checked. Based on the observed colours, the epidermis units were in a suitable condition for use in the assay.

The mean OD value of the three negative control tissues was in the recommended range (0.761). Standard deviation of the viability results for negative control samples was 4.3%.

The positive control treated tissues showed 6.1% viability demonstrating the proper performance of the assay. The standard deviation of the viability results for positive control samples was 0.7%.

The standard deviation of viability values of the three test item-treated tissue samples in the MTT assay was 2.1%.

The mean OD value of the blank samples (acidified isopropanol) was 0.047.

All these parameters met the acceptability criteria, therefore the study was considered to be valid.

Historical control data are presented in Appendix 3.

4.4 Discussion

Following exposure of the EpiSkin™ model to the test item, difenoconazole/fludioxonil/metalaxyl-m/cyclobutifluram FS (A23793B), the mean relative viability was 123.9% compared to the negative control value. This is above the threshold of 50%, therefore under the condition of this assay the test item was considered to be non-irritant to skin. The experiment met the validity criteria, therefore the study was considered to be valid.

5.0 CONCLUSIONS

In conclusion, under the conditions of this *in vitro* EpiSkin™ irritation assay conducted on difenoconazole/fludioxonil/metalaxyl-m/cyclobutifluram FS (A23793B), the results indicate that the test item is non-irritant to skin.

6.0 REFERENCES

1. OECD Guidelines for Testing of Chemicals, Section 4, No. 439, “*In Vitro* Skin Irritation: Reconstructed Human *Epidermis* Test Method”, 14 June 2021
2. Commission Regulation (EC) No 761/2009 of 23 July 2009 amending, for the purpose of its adaptation to technical progress, Regulation (EC) No 440/2008 laying down test methods pursuant to Regulation (EC) No 1907/2006 of the European

Parliament and of the Council on the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), ANNEX III, B.46. IN VITRO SKIN IRRITATION: RECONSTRUCTED HUMAN EPIDERMIS MODEL TEST and amended by Commission Regulation (EU) No 640/2012 of 6 July 2012

3. EpiSkin™ SOP, Version 1.8 (February 2009), ECVAM Skin Irritation Validation Study: Validation of the EpiSkin™ test method 15 min - 42 hours for the prediction of acute skin irritation of chemicals
4. OECD (2015), *Acute Dermal Irritation/Corrosion*, OECD Guideline for the Testing of Chemicals No. 404, OECD, Paris
5. Faller C., Bracher M., Dami N. and Roguet R. (2002) Predictive activity of reconstructed human epidermis equivalents for assessment of skin irritation of cosmetics
6. Mosmann T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunological Methods* 65, 55-62
7. Tinois E., Gaetani Q., Gayraud, B., Dupond D., Rougier A. and Pouradier-Duteil X. (1994) The Episkin model: Successful reconstruction of a human epidermis *in vitro*. In: Rougier, A., Goldberg M., Maibach H.I. Eds. *In Vitro Skin Toxicology*. Mary Ann Liebert, New York, pp. 133-140
8. Hungarian Good Laboratory Practice Regulations: 42/2014. (VIII. 19.) EMMI decree of the Ministry of Human Capacities which corresponds to the OECD GLP, ENV/MC/CHEM(98)17
9. UN (2021), United Nations Globally Harmonized System of Classification and Labelling of Chemicals (GHS), Ninth revised edition, UN New York and Geneva.
10. EC-ECVAM (2009), Statement on the “Performance under UN GHS of three in vitro assays for skin irritation testing and the adaptation of the Reference Chemicals and Defined Accuracy Values of the ECVAM skin irritation Performance Standards”, issued by the ECVAM Scientific Advisory Committee (ESAC30), 9 April 2009
11. EC (2008), REGULATION (EC) No 1272/2008 of the European Parliament and of the Council of 16 December 2008 on classification, labelling and packaging of substances and mixtures, amending and repealing Directives 67/548/EEC and 1999/45/EC, and amending Regulation (EC) No 1907/2006. Official Journal of the European Union L353, 1-1355
12. Guidance on the Application of the CLP Criteria, Guidance to Regulation (EC) No 1272/2008 on classification, labelling and packaging (CLP) of substances and mixtures, Version 5.0, November 2017

7.0 DISTRIBUTION OF THE FINAL REPORT

Sponsor: 1x PDF file, one Word file will be uploaded to collaborative website

Archive: 1x original, bound

SEGREDO INDUSTRIAL

Report Number: 21/245-043B
Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 25 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

TABLES SECTION



SEGREDOS INDUSTRIALIS

Este documento é considerado um Segredo de propriedade da Syngenta Proteção de Cultivos Ltda., constitui-se de SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 26 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

TABLE 1 Optical Density (OD) and the Calculated Non Specific Colour % (NSC_{living}%) of the Additional Control Tissues

Additional control	Optical Density (OD)			NSC% (living)
		Measured	Blank corrected	
Treated with Difenoconazole/Fludioxonil/Metalaxyl-M/ Cyclobutifluram FS (A23793B)	1	0.061	0.015	1.7
	2	0.059	0.012	
	mean	--	0.013	

Notes:

1. Mean blank value was 0.047.
2. Optical density means the mean value of the duplicate wells for each sample (rounded to three decimal places).

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

TABLE 2 Optical Density (OD) and the Calculated Relative Viability % of the Samples

Substance	Optical Density (OD)		Viability (RV%)	Standard Deviation (SD)
	Measured	Blank corrected		
Negative Control: Phosphate buffered saline	1	0.775	0.729	95.8
	2	0.806	0.760	99.8
	3	0.841	0.794	104.4
	mean	--	0.761	100.0
Positive Control: 5% (w/v) SDS solution	1	0.096	0.050	6.5
	2	0.087	0.041	5.3
	3	0.096	0.049	6.4
	mean	--	0.046	6.1
Test Item: Difenoconazole/Fludioxonil/Metalaxyl-M/ Cyclobutifluram FS (A23793B)	1	1.005	0.959	126.0
	2	0.990	0.943	124.0
	3	0.973	0.926	121.7
	mean	--	0.943	123.9

Notes:

1. Mean blank value was 0.047.
2. Optical density means the mean value of the duplicate wells for each sample (rounded to three decimal places).

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

APPENDICES SECTION



CONFIDENTIAL
Property of Syngenta

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

Page 29 of 34

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

APPENDIX 1 Certificate of Analysis



Syngenta Crop Protection, LLC
Analytical and Product Chemistry
Greensboro, NC 27409

Certificate of Analysis

A23793B

Batch ID 1200767 (GP210610)

Test Substance Name:	CGA169374/CGA173506/CGA329351/SYN549522 FS (062.51/049.93/050.05/250.08)
Common Name:	Difenoconazole/Fludioxonil/Metalaxyl-M/Cyclobutifluram FS (062.51/049.93/050.05/250.08)
Material ID:	A23793B
Batch ID:	1200767
Other ID:	GP210610
Source:	Syngenta Crop Protection LLC, 410 Swing Road, Greensboro, NC 27409, US

Chemical Analysis

AI	% w/w	g/L
Difenoconazole	5.45	64.0
Fludioxonil	4.37	51.3
Metalaxyl-M	4.31	50.6
Cyclobutifluram	21.0	247

Identity of the Active Ingredients: Confirmed

Methodology Used for Characterization: LC, mass spectrometry, oscillating density meter.

The Active Ingredient(s) content is within the FAO limits.

Isomer Assay

Analyte	Isomer	% w/w
CGA329351	D-alanine, N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-, Methyl Ester	4.15
CGA351920	L-alanine, N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-, Methyl Ester	0.15

COA Number: USGR210208

Page 1 of 2

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

Physical Analysis

Analyte	Value	Units
---------	-------	-------

Density	1.174	g/cm ³
---------	-------	-------------------

Appearance: red liquid

Storage Temperature: <30°C

Re-certification Date: End of Aug/2024

If stored under the conditions given above, this test substance can be considered stable until the recertification date is reached.

The stability of this test substance will be determined concurrently through reanalysis of material held in inventory under GLP conditions at Syngenta Crop Protection, LLC, Greensboro, NC.

This Certificate of Analysis is summarizing data from a study that has been performed in compliance with Good Laboratory Practices per 40 CFR Part 160. Raw data, documentation, protocols, any amendments to study protocols and reports pertaining to this study are maintained in the Syngenta Crop Protection Archives in Greensboro, NC.

Study Number: USGR210208

Authorization: Sherry Perine

Sherry C Perine

Sherry Perine

Analytical and Product Chemistry Department

Aug 24, 2021

Date

COA Number: USGR210208

Page 2 of 2

SEGREDO INDUSTRIAL

Este documento é considerado de propriedade da Syngenta Proteção de Cultivos Ltda., constitui-se de SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

APPENDIX 2 Copy of the Test Kit Quality Control



NAME

EpiSkin™ Small / Human Epidermis (SM/13)

DESCRIPTION

0.38 cm² reconstructed epidermis of normal human keratinocytes. Cells are grown on a collagen matrix, for 13 days

BATCH : 21-EKIN-042


ORIGIN : Adult donors

USAGE : FOR SCIENTIFIC USE ONLY - PRODUCT OF HUMAN ORIGIN

STORAGE : This product was prepared and packaged using aseptic techniques. Store in an incubator at 37°C, 5% CO₂ with saturated humidity

QUALITY CONTROLS

Control # E211306

HISTOLOGY	Process	Specification	Result
	HES stained paraffin section	Multi-layered, highly differentiated epidermis consisting of organized basal, spinous and granular layers, and a multilayered <i>stratum corneum</i>	Satisfactory
		Number of cell layers ≥ 4	8 cell layers
			
IC50 DETERMINATION	SDS concentration, MTT test.	1.5 mg/mL ≤ IC50 ≤ 3.0 mg/mL	1.9 mg/mL

BIOLOGICAL SAFETY:

On blood of the donors, we have verified the absence of HIV1 and 2 antibodies, hepatitis C antibodies and hepatitis B antigen HBs.

On cells from the donors, we have verified the absence of bacteria, fungus and mycoplasma.

SUGGESTED EXPIRATION DATE:

October 25, 2021

Lyon, October 19, 2021

Certified and released by Michel BATAILLON, Quality Control Manager



Manufactured in accordance to the ISO9001 quality system of Episkin.

The use of this human tissue is strictly limited to *in vitro* testing. All other manipulations of this tissue such as: extraction and maintenance of single cells in culture, use of the tissue for diagnostic or therapeutic purposes and in human subjects, are strictly prohibited.

ISO 9001 Certified

4, rue Alexander Fleming - 69366 Lyon Cedex 07 - France - Tel : +33 (0)4 37 28 72 00 - Fax : +33 (0)4 37 28 72 28
S.A. au capital de 13 608 807 € - 412 127 565 R.C.S. Lyon - NAF : 7211 Z - N° TVA Intracommunautaire FR 46 412 127 565
www.episkin.com



SEGREDOS INDUSTRIALIAIS

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

APPENDIX 3 Historical Control Data

(updated: 25 August 2021)

	Mean value (OD)	Standard deviation (SD)	Minimum value (OD)	Maximum value (OD)	Number of cases
Negative control					
Year 2014-2020	0.811	0.127	0.573	1.381	630
Positive control					
Year 2014-2020	0.062	0.045	0.007	0.354	
Year 2014-2020 viability %	7.6	5.2	0.8	29.7	624

Note: The HC data of the last 7 years are shown in the table. All OD values (measured at 570 ±30 nm) are background corrected values. Before 27 February 2019 the treatment volume was 50µl. After 27 February 2019 the treatment volume was 20µl. After 08 July 2020 the treatment volume is 10µl due to scientific reasons.

Negative control: Phosphate buffered saline (PBS)

Positive control: Sodium dodecyl sulphate (SDS)

SD: Standard deviation

OD: Optical density (absorbance)

Note: All OD values (measured at 570) are background corrected values.

SEGREDO INDUSTRIAL

Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui-se de SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente

APPENDIX 4 GLP Certificate



OGYÉI

Országos Gyógyszerészeti
és Élelmezés-egészségügyi Intézet

Hatósági Ellenőrzési Főosztály

1051 Budapest, Zrínyi utca 3.

Levélírás: 1372 Postafiók 450

Tel.: +36 1 886 9300, Fax: +36 1 886 9460

E-mail: ogyei@ogyei.gov.hu

Web: www.ogyei.gov.hu

Ref. no: OGYÉI/-29520-2/2021

Admin.: Dr. Szaller Zoltán

GOOD LABORATORY PRACTICE (GLP) CERTIFICATE

It is hereby certified that the test facility

Charles River Laboratories Hungary Kft.
H-8200 Veszprém, Szabadságpuszta

is able to carry out

physico-chemical testing, toxicity studies, mutagenicity studies, environmental toxicity studies on aquatic or terrestrial organisms, studies on behaviour in water, soil and air; bio-accumulation, analytical and clinical chemistry, pathology studies, preparation of microscopic tissue sections, reproduction toxicology, in vitro studies, inhalation toxicology, and contract archiving

in compliance with the Principles of GLP (Good Laboratory Practice) and also complies with the corresponding OECD/European Community requirements.

Date of the inspection: 07-11 May 2018.

This certificate is valid up to 11th of May, 2022.

Dr. Lukács
Ferenc
József
Digitalisan aláírta:
Dr. Lukács Ferenc
József
Dátum: 2021.05.06
13:04:14 +02'00'

Dr. Ferenc Lukács
Head of Inspectorate

Note: Translation of the text of the certificate in the header: ("Országos Gyógyszerészeti és Élelmezés-egészségügyi Intézet") - ("National Institute of Pharmacy and Nutrition"); ("Hatósági Ellenőrzési Főosztály") - (Inspectorate Division) and at the signature: ("Digitálisan aláírta") - (Digitally signed); ("Dátum") - ("Date").

SEGREDOES INDUSTRIALIS

Report Number: 21/245-043B Page 34 of 34
Este documento é de propriedade da Syngenta Proteção de Cultivos Ltda., constitui parte do SEGREDO DE NEGÓCIOS e SEGREDO DE INDÚSTRIA, protegidos pelo artigo 195, XI, XII e XIV da Lei N° 9.279/96 e do parágrafo 2º do artigo 9º da Lei 10.603/02.

É terminantemente proibida a divulgação dessas informações e a sua utilização para fins diversos daqueles descritos no parágrafo 2º do artigo 9º da Lei 10.603/02.

Todos os infratores poderão ser processados civil e criminalmente